

**AXIOM A VERSUS NEWHOUSE PHENOMENA FOR BENEDICKS-CARLESON
TOY MODELS
(AXIOME A VERSUS PHÉNOMÈNE DE NEWHOUSE POUR LES MODÈLES
JOUETS DE BENEDICKS-CARLESON)**

CARLOS MATHEUS, CARLOS G. MOREIRA, AND ENRIQUE R. PUJALS

ABSTRACT. We consider a family of planar systems introduced in 1991 by Benedicks and Carleson as a toy model for the dynamics of the so-called Hénon maps. We show that Smale's Axiom A property is C^1 -dense among the systems in this family, despite the existence of C^2 -open subsets (closely related to the so-called Newhouse phenomena) where Smale's Axiom A is violated. In particular, this provides some evidence towards Smale's conjecture that Axiom A is a C^1 -dense property among surface diffeomorphisms.

The basic tools in the proof of this result are: 1) a recent theorem of Moreira saying that stable intersections of dynamical Cantor sets (one of the main obstructions to Axiom A property for surface diffeomorphisms) can be destroyed by C^1 -perturbations; 2) the good geometry of the dynamical critical set (in the sense of Rodríguez-Hertz and Pujals) thanks to the particular form of Benedicks-Carleson toy models.

RÉSUMÉ. Nous considérons une famille de systèmes introduite en 1991 par Benedicks et Carleson comme un modèle jouet pour la dynamique des applications d'Hénon. Nous montrons que l'axiome A de Smale est une propriété C^1 -dense parmi les systèmes dans cette famille, même si nous trouvons aussi des ensembles C^2 -ouverts (liés au phénomène de Newhouse) où l'axiome A de Smale n'est pas satisfait. En particulier, notre résultat soutient la conjecture de Smale selon laquelle l'axiome A est une propriété C^1 -dense parmi les difféomorphismes de surfaces.

Les outils utilisés dans la preuve de notre résultat sont : 1) un théorème récent de Moreira qui dit que les intersections stables des ensembles de Cantor dynamiques (une des obstructions majeures à l'axiome A pour les difféomorphismes de surfaces) peuvent être enlevées par des perturbations C^1 -petites ; 2) la bonne géométrie de l'ensemble de points critiques dynamiques (au sens de Rodríguez-Hertz et Pujals) due à la forme particulière des modèles jouets de Benedicks-Carleson.

1. INTRODUCTION

Uniform hyperbolicity (Smale's Axiom A property) has been a long standing paradigm of complete dynamical description: any dynamical system such that the tangent bundle over its limit set (i.e., the set of accumulation points of all orbits) splits into two complementary subbundles which are uniformly forward (respectively backward) contracted by the tangent map can be completely described from a geometrical and topological point of view.

Date: May 14, 2013.

2000 Mathematics Subject Classification. Primary: 37D40 (Dynamical systems of geometric origin and hyperbolicity); Secondary: 37D20 (Uniformly hyperbolic systems).

Key words and phrases. Axiom A, Newhouse phenomena, Benedicks-Carleson toy models, Hénon maps, dynamical critical points, stable intersections of dynamical Cantor sets, two-dimensional dynamical systems.

Mots-clés. Axiome A, phénomène de Newhouse, modèles jouets de Benedicks-Carleson, applications d'Hénon, points critiques dynamiques, intersections stables des ensembles de Cantor dynamiques, systèmes dynamiques en dimension deux.

Nevertheless, uniform hyperbolicity is a property less universal than it was initially thought: *there are non-empty open sets in the space of dynamics containing only non-hyperbolic systems*. Actually, Newhouse showed that for smooth surface diffeomorphisms, the unfolding of a *homoclinic tangency* (a non transversal intersection of stable and unstable manifolds of a periodic point) generates non-empty open sets of diffeomorphisms whose limit sets are non-hyperbolic (see [N1], [N2], [N3]).

It is important to say that a homoclinic tangency is (locally) easily destroyed by small perturbation of the invariant manifolds. To get open sets of diffeomorphisms with persistent homoclinic tangencies, Newhouse considers certain systems where the homoclinic tangency is associated to an invariant hyperbolic set with large fractal dimension. In particular, he studied the intersection of the local stable and unstable manifolds of a hyperbolic set (for instance, a classical horseshoe), which, roughly speaking, can be visualized as a product of two Cantor sets whose thicknesses are large. Newhouse's construction depends on how this fractal invariant varies with perturbations of the dynamics, and actually this is the main reason that his construction works in the C^2 -topology. In fact, Newhouse argument is based on the continuous dependence of the thickness with respect to C^2 perturbations. A similar construction in the C^1 -topology leading to same phenomena is unknown (indeed, some results in the *opposite* direction can be found in [U] and [M]). In this setting, denoting by $\text{Diff}^r(M^n)$ the set of C^r -diffeomorphisms of a compact n -dimensional manifold M^n (without boundary), it was *implicitly* conjectured by Smale (cf. [Sm], Problems (6.10), item (a), at page 779) that

Axiom A surface diffeomorphisms are C^1 open and dense in $\text{Diff}^1(M^2)$.

This question is explicitly called *Smale's conjecture* in [ABCD].

In the present paper, we consider a special set of maps acting on a two dimensional rectangle, firstly introduced by Benedicks and Carleson as a *toy model* for the so-called *Hénon maps*. For this special type of systems, we show that, if one deals with C^2 -topology, there are non-empty open sets of diffeomorphisms which are not hyperbolic, while in the C^1 -topology, the Axiom A property is open and dense.

Before proceeding further, let us briefly recall some features of Hénon maps and Benedicks-Carleson toy models.

A typical family where the Newhouse's phenomena hold is the so called Hénon maps. In fact, it was proved in [U2] that, for certain parameter of this family, the unfolding of a tangency leads to a non-empty open set of non-hyperbolic diffeomorphisms.

On the other hand, numerical simulations indicate that the attractor of the Hénon map (i.e., the closure of the unstable manifold of its fixed saddle point) has the structure of the product of a line segment and a Cantor set with small dimension (when a certain parameter b is close to zero). Although it is a great oversimplification (and many of the later difficulties on the analysis of Hénon attractors arise because of the roughness of such approximation), this idea gives a very good understanding of the geometry of the Hénon map. As a guide to what follows, it is worth to point out that Benedicks and Carleson [BC, Section 3, p. 89] have constructed a model where the point moves on a pure product space $(-1, 1) \times \mathcal{K}$ where \mathcal{K} is the Cantor set obtained by repeated iteration of the division proportions $(b, 1 - 2b, b)$ (i.e., $\mathcal{K} = \bigcap_{n \geq 0} A^{-n}([0, b] \cup [1 - b, 1])$)

where $A|_{[0,b]}(x) = x/b$ and $A|_{[1-b,1]}(x) = (b - x)/b$, and the dynamics on $(-1, 1)$ is given by a family of quadratic maps: in fact, the dynamical system on $(-1, 1)$ acts as a movement on a fan of lines, where each line has its own x -evolution, while it is contracted in the y -direction (see Figure 1).

More precisely, consider a one parameter family $\{f(x, y)\}_{y \in [0,1]}$ (here x is the variable and y is the parameter) such that, for each fixed parameter $y \in [0, 1]$,

$$f(\cdot, y) : [-1, 1] \rightarrow [-1, 1]$$

is a C^r -unimodal map (with respect to the variable x) verifying that 0 is the critical point and $f(0, y)$ is the maximum value of $f(\cdot, y)$ for all $y \in [0, 1]$. We denote by \mathcal{U}^r the set of families of C^r -unimodal maps satisfying the conditions stated above.

Let $k : [0, a] \cup [b, 1] \rightarrow [0, 1]$ be a C^r function such that $k(0) = 0 = k(1)$, $k(a) = 1 = k(b)$ and $|k'| > \gamma > 1$. Put

$$K(x, y) = \begin{cases} K_+(y) & \text{if } x > 0, \\ K_-(y) & \text{if } x < 0, \end{cases}$$

where $K_+ = (k_{/[0,a]})^{-1}$, $K_- = (k_{/[b,1]})^{-1}$.

The bulk of this article is the study of the dynamics of Benedicks-Carleson toy models $F : ([-1, 1] \setminus \{0\}) \times [0, 1] \rightarrow [-1, 1] \times [0, 1]$ given by

$$(1) \quad F(x, y) = (f(x, y), K(x, y)) = (f(x, y), K_{\text{sgn}(x)}(y)).$$




FIGURE 1. Dynamics of $F(x, y) = (1 - a(y)x^2, K_{\text{sgn}(x)}(y))$ with $a : [0, 1] \rightarrow (0, 2]$.

We denote by \mathcal{D}^r the set of such maps F (with $f(\cdot, y) \in \mathcal{U}^r$ and $k \in C^r$) endowed with the C^r -topology. Since the line $\{x = 0\}$ is a discontinuity line of any $F \in \mathcal{D}^r$, the maps F are C^r -diffeomorphisms only on $([-1, 1] - \{0\}) \times [0, 1]$, and we will explain in Section 2 the exact definition of the C^r -topology. Although this is not specially hard to do, we prefer to postpone it (where we also revise the notion of hyperbolic sets of F) to avoid the appearance of unnecessary technicalities in this introductory section.

At this point, we are ready to state our main results:

Theorem A. *For $r \geq 2$, there exists a non-empty open set $\mathcal{N} \subset \mathcal{D}^r$ such that no $F \in \mathcal{N}$ is Axiom A. Moreover, there exists a residual set $\mathcal{R} \subset \mathcal{N}$ such that any $F \in \mathcal{R}$ has infinitely many periodic sinks.*

On the other hand, in the C^1 -topology, the opposite statement holds:

Theorem B. *There exists an open and dense set $\mathcal{V} \subset \mathcal{D}^1$ such that every $F \in \mathcal{V}$ is Axiom A.*

Concerning the proof of these results, a fundamental role will be played by certain points in the line $\{x = 0\}$:

Definition 1. Given $F \in \mathcal{D}^r$, consider $k : [0, a] \cup [b, 1] \rightarrow [0, 1]$ the Cantor map related to F and denote by \mathcal{K}_0 the Cantor set induced by k . For any $y \in \mathcal{K}_0$, we call

$$c_y^\pm = (0^\pm, y)$$

a critical point of F .

From the technical point of view, it is important to introduce the points $c_y^\pm = (0^\pm, y)$ because we can extend F to them (via the formula $F(0^\pm, y) = (f(0, y), K_\pm(y))$), so that F becomes defined on a compact set. Of course, we have to pay the price that this extension of F is no longer continuous. In particular, let us make a few comments about the orbits and the meaning of the non-wandering set $\Omega(F)$ of this extension of F . While the orbits do not intersect the line $\{x = 0\}$, we have nothing to say. On the other hand, when the iterate $(0, y) = F(z, w)$ of a point $(z, w) \in ([-1, 1] - \{0\}) \times [0, 1]$ hits the line $\{x = 0\}$, we will consider that *both* points $(0^\pm, y)$ make part of the orbit of (z, w) . Finally, we say that a point (z, w) is non-wandering when any neighborhood U of (z, w) has some iterate $F^n(U)$ such that $F^n(U) \cap U \neq \emptyset$. Here, a small neighborhood of a point $(z, w) \in ([-1, 1] - \{0\}) \times [0, 1]$ is a small standard (Euclidean) neighborhood, while a small neighborhood of the point $(0^+, y)$, resp., $(0^-, y)$, is a “half-neighborhood” obtained from the intersection of $[0, 1] \times [0, 1]$, resp. $[-1, 0] \times [0, 1]$, with a small standard (Euclidean) neighborhood of $(0, y)$.

The relevance of the concept of critical point becomes clear from the following simple (but conceptually important) remark:

Remark 1.1. *It follows from the definition that, if $c_y^\pm \in \Omega(F)$ and c_y^\pm is not a periodic sink, then $\Omega(F)$ is not hyperbolic in the sense of Definition 3 below. This fact should be compared to the notion of dynamical critical points of [PRH] and its role as the obstruction to the presence of hyperbolicity/domination in dissipative compact invariant sets of surface diffeomorphisms.*

Closing this introduction, we give the organization of the paper:

- In Section 3, we follow the same ideas of Newhouse to construct a C^2 -open set \mathcal{N} where the critical points can not be removed from the limit set, so that the proof of Theorem A can be derived from the combination of this fact and Remark 1.1.
- In Section 4, the proof of Theorem B is presented. Morally speaking, our basic idea is inspired by a proof of Jakobson’s theorem [J] (of C^1 -density of hyperbolicity among unimodal maps of the interval) along the lines sketched in the book of de Melo and van Strien [dMvS]: namely, in the one-dimensional setting, one combines Mañé’s theorem [M1] (giving the hyperbolicity of compact invariant sets far away from critical points of a C^2 Kupka-Smale interval map) with an appropriate C^1 -perturbation to force the critical point to fall into the basin of a periodic sink. In our two-dimensional setting, we start by showing that the points of the limit set staying away from the critical line $\{x = 0\}$ belong to a hyperbolic set; this is done by proving that any compact set disjoint from the critical line exhibits a dominated splitting and then by using Theorem B in [PS1] (which is the two-dimensional generalization of Mañé’s theorem [M1]) to conclude hyperbolicity. Next, we exploit a recent theorem of Moreira [M] about the non-existence of C^1 -stable intersections of Cantor sets plus the geometry of the maps $F \in \mathcal{D}^1$ to prove a dichotomy for the critical points of a generic F : either critical points fall into the basins of a finite number of periodic sinks or they return to some small neighborhood of the critical line. Finally, we prove the critical points returning close enough to the critical line can be absorbed by the basins of a finite number of periodic

sinks after a C^1 -perturbation; thus, we conclude that the limit set of a generic $F \in \mathcal{D}^1$ is the union of an hyperbolic set with a finite number of periodic sinks, i.e., a generic $F \in \mathcal{D}^1$ is Axiom A.

Acknowledgements. The authors are thankful to IMPA, Collège de France and Institut Mittag-Leffler (and their staff) for the excellent ambient during the preparation of this manuscript. Also, we are grateful to Sylvain Crovisier for several discussions (who helped to clarify the arguments below). Moreover, we would like to acknowledge Sylvain Crovisier and Jean-Christophe Yoccoz for their interest in this work and their constant support.

2. PRELIMINARIES

In this (very) short section, we quickly review a few technical notions appearing in the statements of Theorems A and B.

Definition 2. Given $s \geq r \geq 1$ integers and $F, \tilde{F} \in \mathcal{D}^s$, consider $\{f(\cdot, y)\}_{y \in [0,1]}$ and $k : [0, a] \cup [b, 1] \rightarrow [0, 1]$, respectively $\{\tilde{f}(\cdot, y)\}_{y \in [0,1]}$ and $\tilde{k} : [0, \tilde{a}] \cup [\tilde{b}, 1] \rightarrow [0, 1]$ (the functions associated to F , respectively \tilde{F}).

We say that F and \tilde{F} are C^r -close if the one parameter families $\{f(\cdot, y)\}_{y \in [0,1]}$ and $\{\tilde{f}(\cdot, y)\}_{y \in [0,1]}$ are C^r -close in the usual manner, a is close to \tilde{a} , b is close to \tilde{b} , and k is C^r -close to \tilde{k} in the sense that they admit C^s -extensions to $[0, \max\{a, \tilde{a}\}] \cup [\min\{b, \tilde{b}\}, 1]$ which are C^r -close.

Definition 3. A set Λ is called hyperbolic for $F \in \mathcal{D}^r$ if it is compact, F -invariant and there exist a decomposition $\mathbb{R}^2 = E^s \oplus E^u$ invariant under DF and some constants $C > 0$, $0 < \lambda < 1$ such that

$$|DF_{/E^s(x)}^n| \leq C\lambda^n \text{ and } |DF_{/E^u(x)}^{-n}| \leq C\lambda^n \quad \forall x \in \Lambda, \quad n \in \mathbb{N}.$$

Here, it is worth to point out that we do not require the dimensions of E^s and E^u to be constant on Λ (contrary to some places in the literature). In particular, a hyperbolic set in our context is always the union of a saddle-type set (i.e., a hyperbolic set where $\dim(E^s) = \dim(E^u) = 1$) disjoint from the set of critical points and finitely many periodic sinks.

We say that $F \in \mathcal{D}^r$ is Axiom A if the non-wandering set is hyperbolic and it is the closure of the periodic points. In the sequel, $\Omega(F)$ denotes the non-wandering set (as defined in the paragraph right after Definition 1 above).

3. PROOF OF THEOREM A

The strategy is similar to the arguments of [N1] (see also [PT]).

Given $0 < t < 1$ and $m \geq m_0 = m_0(t)$ (where $m_0(t)$ is a large integer to be chosen later), we define $\delta_m := 1/(2^m - 1)$, $\epsilon_m := \sin(\pi\delta_m/2)$ and we select a parameter ρ_m such that $1 - \cos(\pi\delta_m) < t\rho_m/2 < 1 - \cos(\pi(1 - \delta_m)/2^{m-1})$ (e.g., $\rho_m := 2(1 - \cos(3\pi\delta_m/2))/t$ works for $m_0(t)$ sufficiently large). Next, we take $\mu_m : [0, 1] \rightarrow [0, 1]$ a C^2 -map such that $\mu_m(y) = \mu_m(1 - y)$ and $\mu_m(y) = 1 - \sqrt{1 - \rho_m y/2}$ for every $y \in [0, \frac{t}{2}]$ and we define

$$(2) \quad F^t(x, y) = (f_{\epsilon_m}(x, y), K^t(x, y)),$$

with

$$K^t(x, y) := \begin{cases} (k_{/[0, \frac{t}{2}]}^t)^{-1}(y) & \text{if } x > 0, \\ (k_{/[1 - \frac{t}{2}, 1]}^t)^{-1}(y) & \text{if } x < 0, \end{cases}$$

where k^t is the map

$$k^t(y) = \begin{cases} 2y/t & \text{if } 0 \leq y \leq t/2, \\ 2(1-y)/t & \text{if } 1 - \frac{t}{2} \leq y \leq 1, \end{cases}$$

and $f_{\epsilon_m}(x, y)$ is a C^2 family of unimodal maps such that

$$f_{\epsilon_m}(x, y) = \begin{cases} 1 - 2x^2 & \text{if } |x| \geq \epsilon_m, \\ 1 - \mu_m(y) & \text{at } x = 0. \end{cases}$$

Also, let $\mathcal{K}_0 = \mathcal{K}_0^t := \bigcap_{n \in \mathbb{N}} (k^t)^{-n}([0, \frac{t}{2}] \cup [1 - \frac{t}{2}, 1])$ be the Cantor set induced by $k = k^t$. See Figure 2.




FIGURE 2. Dynamics of K^t .

To simplify the exposition, firstly we consider the proof of Theorem A only for maps $F = F^t$ of the form (2). Then, at the end of this section, we explain how the general case follows from the previous one.

We begin by recalling some classical facts about dynamically defined Cantor sets and their thicknesses. For a more detailed explanation, see [PT].

Definition 4. We say that a Cantor set $\mathcal{K} \subset \mathbb{R}$ is dynamically defined if it is the maximal invariant set of a $C^{1+\alpha}$ expanding map with respect to a given Markov partition.

Definition 5. A gap (resp. bounded gap) of a Cantor set \mathcal{K} is a connected component (resp., bounded connected component) of $\mathbb{R} - \mathcal{K}$. Given U a bounded gap of \mathcal{K} and $u \in \partial U$, we call the bridge C of \mathcal{K} at u to the maximal interval such that $u \in \partial C$ and C contains no point of a gap U' with $|U'| \geq |U|$. The thickness of \mathcal{K} at u is $\tau(\mathcal{K}, u) = |C|/|U|$ and the thickness $\tau(\mathcal{K})$ of \mathcal{K} is the infimum over $\tau(\mathcal{K}, u)$ for all boundary points u of bounded gaps.

Remark 3.1. For the Cantor sets \mathcal{K}_0^t induced by the maps k^t above, it is not hard to see that $0 < \tau(\mathcal{K}_0^t) = t/2(1-t) < \infty$.

Remark 3.2. *The quadratic map $f_2(x) := 1 - 2x^2$ has arbitrarily thick dynamically defined Cantor sets. In fact, using the fact that $1 - 2x^2$ is conjugated to the complete tent map*

$$T_2(x) := \begin{cases} 2x & \text{if } 0 \leq x \leq 1/2, \\ 2 - 2x & \text{if } 1/2 \leq x \leq 1, \end{cases}$$

via the explicit conjugation $h(x) = -\cos(\pi x)$, we can exhibit thick Cantor sets as follows. Denote by $\tilde{I}_2^{(m)} := [h(2\delta_m), h((1 - \delta_m)/2^{m-2})]$ and put $\tilde{I}_i^{(m)} := f_2(\tilde{I}_{i-1}^{(m)})$ for $i = 3, \dots, m$. As it is explained in Section 2 of Chapter 6 of Palis-Takens book [PT], from the explicit nature of the conjugation h and the fact that the intervals $h^{-1}(\tilde{I}_2^{(m)}), \dots, h^{-1}(\tilde{I}_m^{(m)})$ form a Markov partition of a dynamically defined Cantor set \mathcal{K}_m of thickness $\tau(\mathcal{K}_m) = 2^{m-1} - 3$ associated to the tent map $T_2(x)$, it is possible to check that $\tilde{\mathcal{K}}_m := h(\mathcal{K}_m)$ are dynamically defined Cantor sets associated to f_2 (and Markov partition $\tilde{I}_2^{(m)}, \dots, \tilde{I}_m^{(m)}$) such that $\tau(\tilde{\mathcal{K}}_m) \rightarrow \infty$ (as $m \rightarrow \infty$).

Remark 3.3. *Let $\mathcal{K}(\psi)$ be the dynamically defined Cantor set associated to a $C^{1+\alpha}$ expanding map ψ . If ϕ is $C^{1+\alpha}$ -close to ψ , then the thickness of $\mathcal{K}(\phi)$ is close to the thickness of $\mathcal{K}(\psi)$. In other words, the thickness of dynamically defined Cantor sets \mathcal{K} depend continuously on \mathcal{K} (with respect to the $C^{1+\alpha}$ -topology). See [PT].*

Now we state Newhouse's gap lemma ensuring that two linked Cantor sets with large thicknesses should intersect somewhere:

Lemma 3.1 (Gap Lemma [N1]). *Given two Cantor sets \mathcal{K}_1 and \mathcal{K}_2 of \mathbb{R} such that*

$$\tau(\mathcal{K}_1)\tau(\mathcal{K}_2) > 1,$$

then one of the following possibilities occurs:

- \mathcal{K}_1 is contained in a gap of \mathcal{K}_2 ;
- \mathcal{K}_2 is contained in a gap of \mathcal{K}_1 ;
- $\mathcal{K}_1 \cap \mathcal{K}_2 \neq \emptyset$.

For later reference, we recall the following definition:

Definition 6. *We say that \mathcal{K}_1 and \mathcal{K}_2 are linked if their convex hulls I_1 and I_2 are linked in the sense that the interior of I_2 contains exactly one boundary point of I_1 and vice-versa.*

Observe that the “linked” property is robust by perturbations. After these preliminaries, we can complete the discussion of this section as follows.

End of the proof of Theorem A:

We observe that, since $F = F^t$ is the product map $F^t(x, y) = (1 - 2x^2, K_{sgn(x)}^t(y))$ at the region $([-1, \epsilon_m] \cup [\epsilon_m, 1]) \times [0, 1]$, it follows that $\Lambda_{\epsilon_m} := \tilde{\mathcal{K}}_m \times \mathcal{K}_0^t$ is a hyperbolic set of F^t . Moreover, the stable lamination $W^s(\Lambda_{\epsilon_m})$ is composed by vertical lines passing through $\tilde{\mathcal{K}}_m \times \{0\}$ and the unstable lamination $W^u(\Lambda_{\epsilon_m})$ is composed by horizontal lines passing through $\{0\} \times \mathcal{K}_0^t$. We divide the construction of \mathcal{N} into three steps.

Step 1: From Remarks 3.1 and 3.2, given $0 < t < 1$, we can choose $m_0(t) \in \mathbb{N}$ large such that, for every $m \geq m_0(t)$, it holds

$$\tau(\tilde{\mathcal{K}}_m)\tau(\mathcal{K}_0^t) > 2.$$

Step 2: Consider the following line segment:

$$L^+ := F^2(\{0^+\} \times [0, t/2]) = \{(1 - 2(1 - \mu_m(y))^2, \frac{t^2}{4}y)\}_{y \in [0, t/2]} = \{(-1 + \rho_m y, \frac{t^2}{4}y)\}_{y \in [0, t/2]}.$$

In the sequel, L^+ plays the role of a *line of tangencies*: more precisely, we introduce

$$\tilde{\mathcal{K}}^s = (f_2^{-1}(\tilde{I}_2^{(m)} \cap \tilde{\mathcal{K}}_m) \times [0, 1]) \cap L^+, \quad \tilde{\mathcal{K}}^u = F^2(\{0^+\} \times (\mathcal{K}_0^t \cap [0, t/2])) = W_{loc}^u(\Lambda_{\epsilon_m}) \cap L^+.$$

Here, we used the fact that the critical point 0 of f_{ϵ_m} belongs to the interval $\tilde{I}_m^{(m)}$ of the Markov partition of the dynamical Cantor set $\tilde{\mathcal{K}}_m$ (and, thus, the preimages of 0 accumulate $\tilde{\mathcal{K}}_m$) to get that the critical points c_y^\pm belong to $W^u(\Lambda_{\epsilon_m})$.

We claim that $\tilde{\mathcal{K}}^s \cap \tilde{\mathcal{K}}^u \neq \emptyset$. In fact, since the straight line segment L^+ is transversal to both horizontal and vertical foliations, and L^+ is naturally identified with the interval $[0, t/2]$ of \mathbb{R} via $L^+ \ni (-1 + \rho_m y, \frac{t^2}{4} y) \mapsto y \in \mathbb{R}$, we obtain that $\tau(\tilde{\mathcal{K}}^s) \geq \tau(\tilde{\mathcal{K}}_m)/2$ (as the derivative of f_2 in the interval $[0, t/2]$ is between 1 and 2) and $\tau(\tilde{\mathcal{K}}^u) = \tau(\mathcal{K}_0^t)$, so that $\tau(\tilde{\mathcal{K}}^s)\tau(\tilde{\mathcal{K}}^u) > 1$ (by Step 1). Hence, by Newhouse gap lemma 3.1, it suffices to show that $\tilde{\mathcal{K}}^s$ and $\tilde{\mathcal{K}}^u$ are linked. However, it is not hard to see that this follows from our choice of ρ_m . Indeed, from the definitions of $\tilde{\mathcal{K}}^s$ and $\tilde{\mathcal{K}}^u$, we get that $\tilde{\mathcal{K}}^s$ and $\tilde{\mathcal{K}}^u$ are linked if and only if the vertical projection $\bar{\mathcal{K}}^s := f_2^{-1}(\tilde{I}_2^{(m)} \cap \tilde{\mathcal{K}}_m)$ of $\tilde{\mathcal{K}}^s$ is linked to the vertical projection $\bar{\mathcal{K}}^u$ of $\tilde{\mathcal{K}}^u$. On the other hand, the convex hulls of $\bar{\mathcal{K}}^s$ and $\bar{\mathcal{K}}^u$ are linked: more precisely, the convex hull I^s of $\bar{\mathcal{K}}^s$ is $f_2^{-1}(\tilde{I}_2^{(m)}) = [-\cos(\pi\delta_m), -\cos(\pi(1 - \delta_m)/2^{m-1})]$ and the convex hull I^u of $\bar{\mathcal{K}}^u$ is $[0, -1 + t\rho_m/2]$, so that our choice of ρ_m verifying

$$1 - \cos(\pi\delta_m) < t\rho_m/2 < 1 - \cos(\pi(1 - \delta_m)/2^{m-1})$$

implies that I^s and I^u are linked.

Next, we notice that $\tilde{\mathcal{K}}^s \cap \tilde{\mathcal{K}}^u \neq \emptyset$ means that $F^2(c_y^+) \in W_{loc}^s(\Lambda_{\epsilon_m})$ for some critical point $c_y^+ \in W_{loc}^u(\Lambda_{\epsilon_m})$, $y \in \mathcal{K}_0^t$. Since the hyperbolic set Λ_{ϵ_m} is transitive (i.e., it contains dense orbits), it follows that c_y^+ is a non-periodic critical point belonging to the non-wandering set $\Omega(F)$. Therefore, by Remark 1.1, the set $\Omega(F)$ is not hyperbolic.

Step 3: Finally, we claim that any sufficiently small C^2 neighborhood $\mathcal{N} \subset \mathcal{D}^2$ of the map $F = F^t$ constructed above fits the conclusion of the first part of Theorem A. Indeed, this is a consequence of the following known facts for G C^2 -close to F :

- (1) The hyperbolic set Λ_{ϵ_m} has a continuation to a hyperbolic set $\Lambda_{\epsilon_m}(G)$ of G ;
- (2) The Cantor sets \tilde{K}^s and \tilde{K}^u have unique continuations to Cantor sets $\tilde{K}^s(G)$ and $\tilde{K}^u(G)$ obtained by intersecting the local stable and unstable laminations of $\Lambda_{\epsilon_m}(G)$ with the *line of tangencies* $L^+(G) = G^2(\{0^+\} \times [0, 1/2])$. Moreover, these Cantor sets are $C^{1+\alpha}$ -close to \tilde{K}^s and \tilde{K}^u respectively in the sense that their vertical projections $\bar{K}^s(G)$ and $\bar{K}^u(G)$ to $\mathbb{R} \times \{0\}$ are $C^{1+\alpha}$ -close to the vertical projections \bar{K}^s and \bar{K}^u of \tilde{K}^s and \tilde{K}^u ;
- (3) Thus, the Cantor sets $\bar{K}^s(G)$ and $\bar{K}^u(G)$ have thicknesses close to the thicknesses of \bar{K}^s and \bar{K}^u respectively; by continuity of the thickness (see Remark 3.3), it follows that $\tau(\bar{K}^u(G))\tau(\bar{K}^s(G)) > 1$;
- (4) From Newhouse gap lemma 3.1 and the fact that $\bar{K}^s(G)$ and $\bar{K}^u(G)$ remain linked, it follows that $\bar{K}^s(G) \cap \bar{K}^u(G) \neq \emptyset$ and, *a fortiori*, $\tilde{K}^s(G) \cap \tilde{K}^u(G) \neq \emptyset$;
- (5) Hence, there are (non-periodic) critical points contained in the non-wandering set of G , and so, by Remark 1.1, it is not hyperbolic.

At this point, it remains only to prove the second part of Theorem A, namely, the existence of a residual set $\mathcal{R} \subset \mathcal{N}$ such that any $F \in \mathcal{R}$ has infinitely many sinks.

Let $\mathcal{N}_n \subset \mathcal{N}$ be the (open) subset of maps $F \in \mathcal{N}$ with n attracting periodic orbits (at least) disjoint from the critical line $\{x = 0\}$ and $\mathcal{R} = \bigcap_{n \in \mathbb{N}} \mathcal{N}_n$. In this notation, our task is reduced to show the next proposition.

Proposition 3.1. \mathcal{N}_n is dense for every $n \in \mathbb{N}$.

We start our argument with the following notion.

Definition 7. We say that $F \in \mathcal{D}^r$ exhibits a “homoclinic tangency” if there is a hyperbolic periodic point p of saddle-type of F such that

- (1) there exists $c^+ = (0^+, y_p) \in W^u(p)$;
- (2) there exists $k > 0$ such that $F^k(c^+) \in W_{loc}^s(p)$ and $F^j(c^+)$ does not intersect the critical line for $0 < j \leq k$;

The relevance of homoclinic tangencies becomes apparent in the next lemma.

Lemma 3.2. Let $F \in \mathcal{D}^r$ with a homoclinic tangency. Then, there exists $G \in \mathcal{D}^r$ arbitrarily C^r close to F having an attracting periodic orbit disjoint from the critical line $\{x = 0\}$ near the homoclinic tangency.

Proof. From the fact that $c^+ \in W^u(p)$, it follows that there exists $c_n^+ = (0^+, y_n)$ such that $F^{-k_n}(c_n^+) \rightarrow F^k(c^+)$ (for some appropriate sequence k_n) and $c_n^+ \rightarrow c^+ = (0^+, y_\infty)$. Indeed, since the unstable manifold of p consists of horizontal segments (as one can check from the form of the maps $F \in \mathcal{D}^r$), we can apply Palis’ inclination lemma to the vertical segments $\{0^+\} \times (y_\infty - 1/n, y_\infty + 1/n)$ containing c^+ to deduce that, for each $n \in \mathbb{N}$, there are $k_n \in \mathbb{N}$ such that $F^{-k_n}(\{0^+\} \times (y_\infty - 1/n, y_\infty + 1/n))$ intersect a $(1/n)$ -neighborhood of $F^k(c^+)$. Thus, for each $n \in \mathbb{N}$, we can select $c_n^+ = (0^+, y_n) \in \{0^+\} \times (y_\infty - 1/n, y_\infty + 1/n)$ such that $F^{-k_n}(c_n^+)$ belongs to a $(1/n)$ -neighborhood of $F^k(c^+)$. In particular, $c_n^+ \rightarrow c^+$ and $F^{-k_n}(c_n^+) \rightarrow F^k(c^+)$, as desired.

Hence, we can take a C^r small perturbation $H = H_n$ of F such that $H^k(c_n^+) = F^{-k_n}(c_n^+)$ and $H = F$ along the orbit $F^{-j}(c_n^+)$ for $j = 1, \dots, k_n$ (provided that n is large enough). In fact, since $F^k(c^+)$ and the orbit of p don’t meet the critical line, we can fix $\alpha > 0$ such that, for all $n \in \mathbb{N}$, one has $F^j(c_n^+) \in ([-1, -\alpha] \cup [\alpha, 1]) \times [0, 1]$ for $j = 1, \dots, k$ and $F^{-m}(c_n^+) \in ([-1, -\alpha] \cup [\alpha, 1]) \times [0, 1]$ for $m = 1, \dots, k_n + k - 1$. In particular, if we take a smooth bump function $\nu_\alpha : [-1, 1] \rightarrow \mathbb{R}$ with $\nu_\alpha(x) = 0$ if $|x| \geq \alpha$ and $\nu_\alpha(x) = 1$ for $|x| \leq \alpha/2$, and if we set

$$H(x, y) = \left((1 + (X_n - 1)\nu_\alpha(x)) f(x, y), (1 + (Y_n - 1)\nu_\alpha(x)) K_{sgn(x)}(y) \right) \in \mathcal{D}^r,$$

where $X_n := \frac{\pi_1(F^{-k_n-k+1}(c_n^+))}{\pi_1(F(c_n^+))}$, $Y_n := \frac{\pi_2(F^{-k_n-k+1}(c_n^+))}{\pi_2(F(c_n^+))}$, $\pi_1(x, y) = x$ and $\pi_2(x, y) = y$, then we obtain a small C^r -perturbation of $F(x, y) = (f(x, y), K_{sgn(x)}(y))$ because $F^{-k_n}(c_n^+)$ is close to $F^k(c^+)$ and c_n^+ is close to c^+ , so that $F^{-k_n-k+1}(c_n^+)$ and $F(c_n^+)$ are close to $F(c^+)$ and thus X_n and Y_n are close to 1. Moreover, by definition, $H(c_n^+) = F^{-k_n-k+1}(c_n^+)$ and $H = F$ on $\{|x| \geq \alpha\} \times [0, 1]$. Since $F^j(c_n^+)$ belongs to $\{|x| \geq \alpha\} \times [0, 1]$ for $j = -k_n - k + 1, \dots, k$, we obtain that $H^k(c_n^+) = F^{-k_n}(c_n^+)$ and $H = F$ along the orbit $F^{-j}(c_n^+)$ for $j = 1, \dots, k_n$, and, thus, $c_n^+ = H^{k+k_n}(c_n^+)$ is a periodic point of H of period $k + k_n$. Furthermore, using that c_n^+ is a critical point of H , the reader can check that the periodic point c_n^+ is attracting. Of course, this does not complete the argument because the attracting periodic point c_n^+ of $H = H_n$ belongs to the critical line, but this little inconvenience is easily overcome by slightly perturbing $H = H_n$ as follows.

Let us write $H = H_n \in \mathcal{D}^r$ as $H(x, y) = (h(x, y), L_{\text{sgn}(x)}(y))$. Recall that the H -orbit of c_n^+ does not intersect the critical line except for $c_n^+ = (0^+, y_n)$ itself: indeed, using the notation of the previous paragraph, $H^i(c_n^+) \in \{|x| \geq \alpha\} \times [0, 1]$ for all $i = 1, \dots, k + k_n - 1$. Thus, if we denote by $\theta_l = \alpha/2l$, we see that, for all l sufficiently large, we can slightly perturb the unimodal maps $h(x, y_n)$ near $x = 0$ and $h(x, \pi_2(H^{k+k_n-1}(c_n^+)))$ near $x = \pi_1(H^{k+k_n-1}(c_n^+))$ such that the resulting map $G = G_l$ satisfy $G(\theta_l, y_n) = H(c_n^+)$, $G(H^{k+k_n-1}(y)) = (\theta_l, y_n)$ and $G = H$ nearby the points $H^i(c_n^+)$, $i = 1, \dots, k + k_n - 2$. In particular, the resulting map $G = G_l \in \mathcal{D}^r$ has an attracting periodic point of period $k + k_n$ at (θ_l, y_n) whose G -orbit does not meet the critical line. This ends the proof. \square

On the other hand, homoclinic tangencies are frequent inside \mathcal{N} .

Lemma 3.3. *Let $F \in \mathcal{N}$. Then, there exists $G \in \mathcal{D}^r$ close to F exhibiting a homoclinic tangency.*

Proof. This is an immediate consequence of the construction of \mathcal{N} : given $F \in \mathcal{N}$, we can find $x_1, x_2 \in \Lambda_{\epsilon_m}$ and a critical point $c^+ \in W_{loc}^u(x_1)$ such that $F^k(c^+) \in W_{loc}^s(x_2)$. Since the hyperbolic set Λ_{ϵ_m} is transitive, we can find a periodic p in Λ_{ϵ_m} close to x_1 so that some pieces of its local unstable and stable manifolds are close to the corresponding invariant manifolds of x_1 and x_2 (resp.). In this situation, after a proper small perturbation (similar to the ones performed during the proof of Lemma 3.2 above), we can find $G \in \mathcal{D}^r$ C^r -close to F such that $G \in \mathcal{D}^r$ has a homoclinic tangency involving p . \square

Finally, the proof of the desired proposition follows from a direct combination of the two previous lemmas.

Proof of Proposition 3.1. It is proved by induction. Given $G_n \in \mathcal{N}_n$, we can use Lemma 3.3 to find $G_{n+1} \in \mathcal{D}^r$ C^r -close to G_n keeping the same number n of attracting periodic points of G_n and such that G_{n+1} has a homoclinic tangency. By Lemma 3.2, we can unfold this tangency to create a new sink, i.e., we can find $H \in \mathcal{N}_{n+1}$ C^r -close to G_{n+1} . The result follows by Baire's theorem. \square

This completes the proof of Theorem A.

4. PROOF OF THEOREM B

Before giving the proof of Theorem B, we briefly outline the strategy. Given $\epsilon > 0$, let us take $U_\epsilon = ([-1, -\epsilon] \cup [\epsilon, 1]) \times [0, 1]$ and

$$\Lambda_\epsilon = \Omega(F) \cap \bigcap_{n \in \mathbb{Z}} F^n(U_\epsilon).$$

Strategy of the proof.

- (1) For any $\epsilon > 0$, we show that, C^1 -generically, the set Λ_ϵ is composed by a locally maximal hyperbolic set and a finite number of periodic attracting points. This is performed in Subsection 4.1 (see Theorem 4.1 and Corollary 4.1).
- (2) We show that, C^1 -generically, any critical point either it is contained in the basin of attraction of the sinks (of Step 1 above) or it returns to $[-\epsilon, \epsilon] \times [0, 1]$. This is performed in Subsection 4.2.
- (3) Later, we produce a series of C^1 -perturbations (of size proportional to ϵ) in the way to create a finite number of periodic sinks such that their basins contain the critical points that return. This is performed in Subsection 4.3.

- (4) From items 1, 2 and 3, it follows that $\Omega(F) \subset \Lambda_\epsilon \cup \{p_1, \dots, p_k\}$, where each p_i is a periodic attracting point ($i = 1, \dots, k$), and therefore it is concluded that $\Omega(F)$ is hyperbolic (and F is Axiom A).

Convention. As it will become clear later, although our Theorem B concerns \mathcal{D}^1 with the C^1 -topology, sometimes we will need some extra (C^2) regularity of the maps (for instance, this is the case when one tries to apply a fundamental “hyperbolicity criterion for surface maps” of Pujals and Sambarino). Therefore, during this *entire* (last) Section 4, we consider \mathcal{D}^2 equipped with the C^1 -topology. In particular, each time we refer to “a C^1 -generic/typical $F \in \mathcal{D}^2$ ”, we mean a $F \in \mathcal{D}^2$ belonging to an appropriate C^1 -open and C^1 -dense subset of \mathcal{D}^2 .

4.1. Hyperbolicity of Λ_ϵ .

Theorem 4.1. *Let $\epsilon > 0$ be a positive constant. Then, there exists a C^1 -open and dense subset $\mathcal{V}_\epsilon \subset \mathcal{D}^1$ such that, for any $F \in \mathcal{V}_\epsilon$, the set Λ_ϵ contains a finite number of periodic attracting points and the complement of the basin of attraction of them $\hat{\Lambda}_\epsilon$ exhibits a hyperbolic splitting $T\hat{\Lambda}_\epsilon = E^s \oplus E^u$ such that E^s is contracting, E^u is expanding (and, in fact, $E^u = \mathbb{R} \cdot (1, 0)$).*

The proof of this result uses the notion of dominated splitting and Theorem B in [PS1]. Firstly, we revisit the definition of dominated splittings:

Definition 8. *A f -invariant set Λ has a dominated splitting if we can decompose its tangent bundle into two invariant subbundles $T_\Lambda M = E \oplus F$ such that:*

$$(3) \quad \|Df^n_{/E(x)}\| \cdot \|Df^{-n}_{/F(f^n(x))}\| \leq C\lambda^n, \text{ for all } x \in \Lambda, n \geq 0.$$

with $C > 0$ and $0 < \lambda < 1$.

Secondly, we recall that Pujals and Sambarino [PS1] proved that any compact invariant set exhibiting dominated splitting of a generic C^2 surface diffeomorphism is hyperbolic:

Theorem 4.2 ([PS1]). *Let $f \in \text{Diff}^2(M^2)$ be a C^2 -diffeomorphism of a compact surface M^2 and $\Lambda \subset \Omega(f)$ a compact invariant set exhibiting a dominated splitting. Assume that all periodic points in Λ are hyperbolic of saddle type. Then, Λ can be decomposed into a hyperbolic set and a finite number of normally hyperbolic periodic closed curves whose dynamical behaviors are C^2 -conjugated to irrational rotations.*

Let us begin the proof of Theorem 4.1 with some useful notation. Given (x_0, y_0) , we denote by $(x_i, y_i) := F^i(x_0, y_0)$; also, we write the derivative of a map $F(x, y) = (f(x, y), K(x, y))$ of the form (1) as

$$DF = \begin{pmatrix} f_x & f_y \\ 0 & K_y \end{pmatrix}.$$

In particular, it follows that

$$DF^n(x_0, y_0) = \begin{pmatrix} A_n & B_n \\ 0 & D_n \end{pmatrix},$$

where

$$A_n := \prod_{i=0}^{n-1} f_x(x_i, y_i) \quad , \quad D_n := \prod_{i=0}^{n-1} K_y(x_i, y_i),$$

$$B_n = \sum_{j=0}^{n-1} f_y(x_j, y_j) \prod_{i=0}^{j-1} K_y(x_i, y_i) \prod_{i=j+1}^{n-1} f_x(x_i, y_i).$$

In the sequel, we fix two positive constants λ_0, λ_1 such that $\lambda_0 < \lambda_1 < 1$ and

$$|K_y| < \lambda_0.$$

Concerning the proof of Theorem 4.1, we observe that $(1, 0)$ is an invariant direction by DF and, moreover, it is the natural candidate to be the expanding one. Therefore, the existence of a dominated splitting follows once we build up a proper invariant cone field around $(1, 0)$. To perform this task, first we need the next lemma.

Lemma 4.1. *Given $\epsilon > 0$, let $\mathcal{Z}_\epsilon \subset \mathcal{D}^1$ be the C^1 -open and dense subset consisting of $F \in \mathcal{D}^1$ such that, for each $y \in [0, 1]$, the unimodal map $f(\cdot, y)$ has no critical points in $|x| \geq \epsilon$. Then, for any $F \in \mathcal{Z}_\epsilon$, there exist a finite number of attracting periodic points with trajectory in Λ_ϵ and a positive integer $n_0 = n_0(\epsilon)$ such that, for any $(x_0, y_0) \in \Lambda_\epsilon$ outside the basins of attraction of those periodic points, it holds*

$$|A_n| = \prod_{i=0}^{n-1} |f_x(x_i, y_i)| > \lambda_1^n,$$

whenever $n > n_0$.

In order to do not interrupt the flow of ideas, we postpone the proof of the lemma. Assuming momentarily this lemma, we are able to prove Theorem 4.1.

Proof of Theorem 4.1. Let b be a positive constant such that

$$|f_y| < b.$$

Given $F \in \mathcal{Z}_\epsilon$, we can take n_0 the integer provided by Lemma 4.1 and let R_0 be a positive constant¹ such that, for any $m < n_0$ and any point $(x_0, y_0) \in \Lambda_\epsilon$, it holds

$$\prod_{i=0}^m |f_x(x_i, y_i)| > R_0^{-1}.$$

Now, for all $(x_0, y_0) \in \Lambda_\epsilon$ outside the basins of the attracting periodic points of Lemma 4.1, let us bound B_n for $n > n_0$:

$$\begin{aligned} (4) \quad |B_n| &\leq \sum_{j=0}^{n-1} |f_y(x_j, y_j)| \prod_{i=0}^{j-1} |K_y(x_i, y_i)| \prod_{i=j+1}^{n-1} |f_x(x_i, y_i)| \\ &= \sum_{j=0}^{n-1} |f_y(x_j, y_j)| \cdot |D_j| \cdot \frac{|A_n|}{|A_{j+1}|} \\ &< R_0 b |A_n| \frac{1}{1 - \lambda_0} + b |A_n| \sum_{j=n_0}^{n-1} \frac{\lambda_0^j}{\lambda_1^j} \\ &< R_0 b |A_n| \frac{1}{1 - \lambda_0} + b |A_n| \frac{1}{\lambda_1 - \lambda_0}. \end{aligned}$$

Using this estimate, we claim that the cone field $C(\gamma_0) := C(\mathbb{R} \cdot (1, 0), \gamma_0) := \{(\dot{x}, \dot{y}) \in \mathbb{R}^2 : |\dot{y}| \leq \gamma_0 |\dot{x}|\}$ is a forward invariant cone field for sufficiently small $\gamma_0 > 0$. In fact, take $\gamma_0 > 0$ small and let us consider

$$v_n = DF^n(1, \gamma) = (A_n + \gamma B_n, \gamma D_n),$$

¹Such a constant R_0 always exists since $f(x, y) \in \mathcal{U}^1$ is a family of unimodal maps without critical points in $|x| \geq \epsilon$ for $F \in \mathcal{Z}_\epsilon$ and $(x_i, y_i) \in U_\epsilon$ implies $|x_i| \geq \epsilon$.

where $|\gamma| < \gamma_0$. The slope of v_n with respect to $(1, 0)$ is

$$|\text{slope}(v_n, (1, 0))| = \frac{|\gamma D_n|}{|A_n + \gamma B_n|}.$$

Note that the estimate (4) implies

$$|A_n + \gamma B_n| > |A_n| - \gamma_0 |B_n| > |A_n| (1 - \gamma_0 b(R_0 + 1) \cdot (\lambda_1 - \lambda_0)^{-1}).$$

Hence, if γ_0 is small so that

$$1 - \gamma_0 b(R_0 + 1)(\lambda_1 - \lambda_0)^{-1} > \frac{1}{2},$$

using Lemma 4.1, we conclude that

$$\text{slope}(v_n, (1, 0)) < \gamma_0 \frac{2|D_n|}{|A_n|} < 2 \left(\frac{\lambda_0}{\lambda_1} \right)^n \cdot \gamma_0.$$

Thus, assuming n_0 large so that $(\lambda_0/\lambda_1)^{n_0} < 1/4$ and taking $\gamma_1 = 2(\lambda_0/\lambda_1)^{n_0} \gamma_0$, we see that, for any $n > n_0$,

$$DF^n(C(\gamma_0)) \subset C(\gamma_1) \subset C(\gamma_0/2).$$

In other words, $C(\gamma_0)$ is a forward invariant cone field and the existence of a dominated splitting $E^s \oplus \mathbb{R} \cdot (1, 0)$ is guaranteed (over the set $\hat{\Lambda}_\epsilon$ of points outside the basins of the attracting points of Lemma 4.1).

Next, we show that E^s is uniformly contracted: for every $(x_0, y_0) \in \Lambda_\epsilon$, we fix $e_0^{(s)} = (u_0^s, v_0^s) \in E_{(x_0, y_0)}^s$ with $\|e_0^{(s)}\| = 1$ and we put $DF^n(x_0, y_0) \cdot e_0^{(s)} := \pm \lambda_n^s \cdot e_n^{(s)} \in E_{(x_n, y_n)}^s$ where $e_n^{(s)} := (u_n^{(s)}, v_n^{(s)}) \in E_{(x_n, y_n)}^s$ is an unitary vector. Then, we compute the determinant of DF^n :

$$|A_n \cdot D_n| = |\det DF^n| = \frac{|DF^n \cdot (1, 0) \wedge DF^n \cdot e_0^{(s)}|}{|(1, 0) \wedge e_0^{(s)}|} = \frac{|A_n| \cdot |\lambda_n^s| \cdot |v_n^{(s)}|}{|v_0^{(s)}|},$$

where $|u \wedge v|$ denotes the area of the rectangle determined by the vectors u and v . Because the direction E^s does not belong to the cone field $C(\gamma_0)$ and $|v_0^{(s)}| \leq \|e_0^{(s)}\| = 1$, we get

$$|\lambda_n^s| = |D_n| \frac{|v_0^{(s)}|}{|v_n^{(s)}|} \leq \frac{1}{\gamma_0} |D_n|.$$

Since $|D_n| \leq \lambda_0^n$ for all $n \in \mathbb{N}$, this proves that for all $F \in \mathcal{Z} \subset \mathcal{D}^1$ of the form (1) such that $|K_y| < \lambda_0$ and for any $\lambda_0 < \lambda_1 < 1$, the set Λ_ϵ is the union of a finite number of sinks and a set $\hat{\Lambda}_\epsilon$ exhibiting a dominated decomposition $E^s \oplus F$ where E^s is contracting (after n_0 iterates) and $F = (1, 0) \cdot \mathbb{R}$ satisfies $DF^n(1, 0) = (A_n, 0)$ where $|A_n| > \lambda_1^n$ (for $n > n_0$).

At this stage, the proof of Theorem 4.1 is reduced to show that there exists a C^1 -dense subset of $F \in \mathcal{D}^2 \cap \mathcal{Z}_\epsilon$ such that all periodic points in $\hat{\Lambda}_\epsilon$ are hyperbolic of saddle type. Indeed, this is true because it is immediate that there are no periodic closed curves inside Λ_ϵ whose dynamical behavior are conjugated to irrational rotations², so that, by Pujals-Sambarino Theorem 4.2, a dominated splitting over $\hat{\Lambda}_\epsilon$ is a hyperbolic splitting whenever all periodic points in $\hat{\Lambda}_\epsilon$ are hyperbolic of saddle type.

²Because in this case $F = \mathbb{R} \cdot (1, 0)$ should be a tangent line of such closed curve C at some point. Combining this fact with the minimality of the dynamics on C and the continuity of dominated splitting (besides the invariance of $\mathbb{R} \cdot (1, 0)$), we obtain that the whole curve C is tangent to the line field F , a contradiction.

However, the C^1 -denseness in \mathcal{D}^1 of $F \in \mathcal{D}^2 \cap \mathcal{Z}_\epsilon$ such that all periodic points in $\hat{\Lambda}_\epsilon$ are hyperbolic of saddle type is a consequence of a simple argument (compare with [PS1, p. 966]): recall that, by the “usual” transversality arguments, all periodic points of a $F \in \mathcal{D}^2 \cap \mathcal{Z}_\epsilon$ in a C^2 -generic (G_δ dense) subset are hyperbolic³; it follows that for such a $F \in \mathcal{D}^2 \cap \mathcal{Z}_\epsilon$, the compact invariant subset $\Lambda_\epsilon^{(0)} := \Lambda_\epsilon - \{p \in \Lambda_\epsilon : p \text{ is a periodic sink}\} \subset \Omega(F)$ only contains hyperbolic periodic points of saddle type. Furthermore, $\Lambda_\epsilon^{(0)}$ admits a dominated splitting (since $\Lambda_\epsilon^{(0)} \subset \hat{\Lambda}_\epsilon$). Thus, we obtain from Theorem 4.2 that $\Lambda_\epsilon^{(0)}$ is a hyperbolic set. We claim that $P_\epsilon(F) := \Lambda_\epsilon - \Lambda_\epsilon^{(0)}$ is finite (so that $\Lambda_\epsilon^{(0)} = \hat{\Lambda}_\epsilon$ and, *a fortiori*, all periodic points of $\hat{\Lambda}_\epsilon$ are hyperbolic of saddle-type). Indeed, if $\#P_\epsilon(F) = \infty$, we have $\emptyset \neq \overline{P_\epsilon(F)} - P_\epsilon(F) \subset \Lambda_\epsilon^{(0)}$. However, since $\Lambda_\epsilon^{(0)}$ is hyperbolic, we can select a compact neighborhood U of Λ_ϵ such that the maximal invariant of U is hyperbolic. Thus, we get that, up to removing a finite number of periodic sinks, $P_\epsilon(F) \subset U$, a contradiction with the hyperbolicity of the maximal invariant subset of U . This completes the proof of Theorem 4.1. \square

Closing the proof of the hyperbolicity of Λ_ϵ , we prove the statement of Lemma 4.1.

Proof of Lemma 4.1. It is enough to apply the following lemma due to Pliss (see [PI], [M2]).

Lemma 4.2 (Pliss). *Given $0 < \gamma_0 < \gamma_1 < 1$ and $a > 0$, there exist $n_0 = n_0(\gamma_0, \gamma_1, a)$ and $l = l(\gamma_0, \gamma_1, a) > 0$ such that, for any sequences of numbers $\{a_i\}_{0 \leq i \leq n}$ with $n_0 < n$, $a^{-1} < a_i < a$ and $\prod_{i=0}^n a_i < \gamma_0^n$, there are $1 \leq n_1 < n_2 < \dots < n_r \leq n$ with $r > ln$ and such that*

$$\prod_{i=n_j}^k a_i < \gamma_1^{k-n_j} \quad \text{for all } n_j \leq k \leq n.$$

In fact, let us consider the set of points $(z, w) \in \Lambda_\epsilon$ such that

$$(5) \quad \liminf_{n \rightarrow \infty} \frac{1}{n} \log |A_n(z, w)| < \log \sqrt{\lambda_1}.$$

Since $F \in \mathcal{Z}_\epsilon$ and for any $(z_i, w_i) = F^i(z, w) \in \Lambda_\epsilon$, it holds $|z_i| \geq \epsilon$, we have that the numbers $a_i = f_x(z_i, w_i)$ are uniformly bounded away from zero and infinity (i.e., $0 < a(\epsilon)^{-1} < a_i < a(\epsilon) < \infty$), and, thus, we can use Lemma 4.2 twice to obtain that there exists a subsequence of forward iterates of (z, w) accumulating on some point (x_0, y_0) which has a subsequence of forward iterates

$$\{(x_{n_j}, y_{n_j})\}_{j>0} = \{F^{n_j}(x_0, y_0)\}$$

such that any (x_{n_j}, y_{n_j}) satisfies

$$|A_n(x_{n_j}, y_{n_j})| < \sqrt{\lambda_1^n}, \quad \forall n > 0.$$

Using the same type of calculation of estimative (4), we get, for any $j > 0$,

$$\prod_{i=0}^n \|DF(x_{i+n_j}, y_{i+n_j})\| < (1 + b(\sqrt{\lambda_1} - \lambda_0)^{-1}) \sqrt{\lambda_1^n}, \quad \forall n > 0 \text{ large.}$$

By standard arguments it follows that, for any $\sqrt{\lambda_1} < \lambda_2 < 1$, there exists $\gamma = \gamma(\lambda_1, \lambda_2)$ such that

$$F^n(B_\gamma(x_{n_j}, y_{n_j})) \subset B_{\lambda_2^n \gamma}(F^n(x_{n_j}, y_{n_j}))$$

³In our context of F of the form (1), the derivative DF is an upper triangular matrix whose diagonal entries are the x -derivative of $f(x, y)$ and the y -derivative of $K_\pm(y)$. In particular, given any non-hyperbolic periodic point of F , we can slightly perturb K_\pm to slightly change both the trace and the determinant of DF to get an hyperbolic periodic point. Therefore, the set of $F \in \mathcal{D}^2$ whose periodic points of period $\leq n$ are hyperbolic are C^2 -open and dense, and, hence, the set of $F \in \mathcal{D}^1$ whose periodic points are all hyperbolic form a C^2 G_δ -dense.

for all $j, n > 0$ large. Taking q_0 an accumulation point of $\{(x_{n_j}, y_{n_j})\}$, it is not hard to see that

$$F^j(B_\gamma(q_0)) \subset B_{\lambda_2^j \gamma}(F^j(q_0))$$

for any $j > 0$ large and there exists a positive integer $m = m(q_0)$ such that

$$F^m(B_\gamma(q_0)) \subset B_{\lambda_2^m \gamma}(q_0).$$

Therefore, it follows that:

- (1) there is an unique attracting periodic point⁴ p_0 inside $B_\gamma(q_0)$,
- (2) the neighborhood $B_\gamma(q_0)$ is contained in the basin of attraction of p_0 ,
- (3) the point (x_0, y_0) and the initial point (z, w) verifying (5) belong to the basin of attraction p_0 ;

Since the number of attracting periodic point with local basin of attraction with radius larger than γ is finite, we conclude that there are a finite number of periodic attracting points whose basins contain the points of Λ_ϵ verifying (5). In other words, Λ_ϵ is the union of finitely many periodic sinks and the subset $\hat{\Lambda}_\epsilon$ of Λ_ϵ consisting of the points violating (5). Note that $\hat{\Lambda}_\epsilon$ is compact because it lies in the complement of the basins of attraction of finitely many sinks. At this point, the proof of the lemma will be complete if we show that there exists $n_0 = n_0(\epsilon) \in \mathbb{N}$ such that $|A_n(z, w)| > \lambda_1^n$ for all $n > n_0$ and $(z, w) \in \hat{\Lambda}_\epsilon$. However, the existence of such an integer n_0 follows from Pliss lemma and the arguments of the previous paragraphs. Indeed, the non-existence of n_0 would imply that there exists a sequence $(z_i, w_i) \in \hat{\Lambda}_\epsilon$ and $n_i \in \mathbb{N}$, $n_i \rightarrow \infty$, such that $|A_{n_i}(z_i, w_i)| \leq \lambda_1^{n_i}$. Using Pliss' lemma (as in the previous paragraphs), we would be able to extract a subsequence of (z_i, w_i) accumulating in some point (z_∞, w_∞) verifying (5). Of course, this is a contradiction because $\hat{\Lambda}_\epsilon$ is compact (and thus $(z_\infty, w_\infty) \in \hat{\Lambda}_\epsilon$ and this concludes the proof of the lemma. \square

For later use, we observe that the hyperbolic sets Λ_ϵ can be assumed to be locally maximal. This follows from the next claim (compare with [A]).

Claim 1. *There exists a locally maximal hyperbolic set $\Lambda_\epsilon \subset \tilde{\Lambda}_\epsilon \subset U_{\epsilon/2}$. In other words, there exists an open set such $V \subset U_{\epsilon/2}$ containing Λ_ϵ such that $\tilde{\Lambda}_\epsilon := \bigcap_{n \in \mathbb{Z}} F^n(V)$ is a compact invariant hyperbolic set.*

Proof. Indeed, fix $\gamma = \gamma(\epsilon) > 0$ a positive small constant such that the local stable manifold $W_\gamma^s(p)$ of any point $p \in \Lambda_{\epsilon/2}$ is the graph of a real function of the y -coordinate defined over an interval of length $\delta = \delta(\epsilon) > 0$. Next, we take $k = k(\epsilon) > 0$ a large integer so that the lengths of the 2^k intervals $I_1^{(k)}, \dots, I_{2^k}^{(k)}$ of the k th stage of the construction of the Cantor set \mathcal{K}_0 are $< \delta/2$. Note that we can suppose that $W_\gamma^s(p) \subset U_{\epsilon/2}$ for any $p \in \Lambda_{\epsilon/2} \cap U_{3\epsilon/4}$. Now, for each $j = 1, \dots, 2^k$, we consider the stable lamination $\mathcal{F}_{j,\pm}^s = \{W_\gamma^s(p) \cap [-1, 1] \times I_j^{(k)}\}_{p \in \hat{\Lambda}_{\epsilon/2} \cap U_{3\epsilon/4}}$. Given $\ell \in \mathcal{F}_{j,-}^s$, resp. $\ell \in \mathcal{F}_{j,+}^s$, we denote by $R_{j,-}^{(k)}(\ell)$, resp. $R_{j,+}^{(k)}(\ell)$, the rectangle delimited by the four lines $\{-1\} \times [0, 1]$, $[-1, 1] \times \partial I_j^{(k)}$ and ℓ , resp. $\{+1\} \times [0, 1]$, $[-1, 1] \times \partial I_j^{(k)}$ and ℓ . Given $\ell, \tilde{\ell} \in \mathcal{F}_{j,\pm}^s$, we say that $\ell \prec \tilde{\ell}$ if and only if $R_{j,\pm}^{(k)}(\ell) \subset R_{j,\pm}^{(k)}(\tilde{\ell})$. Observe that \prec is a total order⁵ of $\mathcal{F}_{j,\pm}^s$. Thus, for each $j = 1, \dots, 2^k$, we can define $\ell_{j,\pm} \in \mathcal{F}_{j,\pm}^s$ the outermost stable leaf of $\hat{\Lambda}_{\epsilon/2} \cap U_{3\epsilon/4} \cap [-1, 1] \times I_j^{(k)}$ as the unique leaf of $\mathcal{F}_{j,\pm}^s$ such that $\ell \prec \ell_{j,\pm}$ for

⁴Actually, using that $(x_{n_j}, y_{n_j}) = F^{n_j}(x_0, y_0) \rightarrow q_0$, it can be concluded that q_0 is the periodic point.

⁵Because any two distinct stable leaves are disjoint and $\partial \ell \subset [-1, 1] \times \partial I_j^{(k)}$ for any $\ell \in \mathcal{F}_{j,\pm}^s$.

all $\ell \in \mathcal{F}_{j,\pm}^s$. Consider the family of rectangles $R_{j,\pm}^{(k)}(\epsilon) := R_{j,\pm}^{(k)}(\ell_{j,\pm})$. Finally, let $\tilde{\Lambda}_\epsilon$ be the maximal invariant set associated to this family of rectangles. It follows that $\tilde{\Lambda}_\epsilon$ has local product structure (because $W_{loc}^s(p) \cap W_{loc}^u(q) \in R_{j,\pm}^{(k)}(\epsilon)$ when $p, q \in \tilde{\Lambda}_\epsilon \cap R_{j,\pm}^{(k)}(\epsilon)$) and $\Lambda_\epsilon \subset \tilde{\Lambda}_\epsilon \subset U_{\epsilon/2}$ (because $\Lambda_\epsilon \cap [-1, 1] \times I_j^{(k)} \subset R_{j,\pm}^{(k)} \subset U_{\epsilon/2}$). This proves our claim. \square

Claim 2. *The set $\tilde{\Lambda}_\epsilon$ found in Claim 1 is the maximal invariant set of $U_\epsilon \cup \widetilde{R}_\epsilon$, where $\widetilde{R}_\epsilon = \{R_{j,\pm}^{(k)}(\ell_{j,\pm})\}$ is the family of rectangles introduced in the previous claim.*

Proof. Indeed, given z a point whose orbit $\mathcal{O}(z)$ stays in $U_\epsilon \cup \widetilde{R}_\epsilon$, we note that $z \in \Lambda_{\epsilon/2}$ (since $U_\epsilon \cup \widetilde{R}_\epsilon \subset U_{\epsilon/2}$). On the other hand, we have two possibilities:

- $\mathcal{O}(z) \subset \widetilde{R}_\epsilon$: this means that $z \in \tilde{\Lambda}_\epsilon$;
- there exists $y \in \mathcal{O}(z) - \widetilde{R}_\epsilon$: this means that $y \in (U_\epsilon \cap \Lambda_{\epsilon/2}) - \widetilde{R}_\epsilon$, a contradiction (since, by definition, $U_\epsilon \cap \Lambda_{\epsilon/2} \subset U_{3\epsilon/4} \cap \Lambda_{\epsilon/2} \subset \widetilde{R}_\epsilon$).

In particular, it follows that the positive orbit $\mathcal{O}^+(p)$ of every point $p \notin W^s(\tilde{\Lambda}_\epsilon)$ escapes any sufficiently small neighborhood of $U_\epsilon \cup \widetilde{R}_\epsilon$. In fact, if the positive orbit of a given point p stays forever inside a small neighborhood W of $U_\epsilon \cup \widetilde{R}_\epsilon$, its accumulation points always belong to the maximal invariant set $\Lambda(W)$ of W . However, since the maximal invariant $\tilde{\Lambda}_\epsilon$ of $U_\epsilon \cup \widetilde{R}_\epsilon$ is locally maximal (by Claim 1), $\Lambda(W) = \tilde{\Lambda}_\epsilon$ for any small neighborhood W of $U_\epsilon \cup \widetilde{R}_\epsilon$. Hence, $p \in W^s(\tilde{\Lambda}_\epsilon)$, a contradiction. \square

Before proceeding further, we use a fundamental result of C. G. Moreira to improve the geometry of the isolating neighborhood of $\tilde{\Lambda}_\epsilon$.

Theorem 4.3 ([M]). *Let \mathcal{K} be a C^2 -dynamically defined Cantor set and let $\tilde{\mathcal{K}}$ be a C^1 -dynamically defined Cantor set. Then, there are C^1 -dynamically defined Cantor sets $\widehat{\mathcal{K}}$ arbitrarily C^1 -close to $\tilde{\mathcal{K}}$ such that $\mathcal{K} \cap \widehat{\mathcal{K}} = \emptyset$. In particular, generically in the C^1 -topology, a pair of C^1 -dynamically defined Cantor sets are disjoint and the arithmetic difference of a C^1 generic pair of C^1 -dynamically defined Cantor sets has empty interior (so that it is also a Cantor set).*

More precisely, combining our Theorem 4.1 with this theorem, we have the following consequence:

Corollary 4.1. *Fix $\epsilon > 0$. Then, for F in a C^1 -open and dense subset of \mathcal{D}^1 , the maximal invariant set Λ_ϵ of U_ϵ is a locally maximal hyperbolic set such that $\text{int}(U_\epsilon)$ is an isolating neighborhood of Λ_ϵ .*

Proof. Let $F \in \mathcal{V}_{\epsilon/2} \cap \mathcal{Z}_\epsilon \cap \mathcal{D}^2$ where $\mathcal{V}_{\epsilon/2}$ is the C^1 -open and dense subset of \mathcal{D}^1 verifying Theorem 4.1. We consider a finite Markov partition $\{P_i\}_{i=1}^M$ of $\widetilde{\Lambda}_{\epsilon/2}$ with small diameter. We take $p_i \in P_i \cap \widetilde{\Lambda}_{\epsilon/2}$ and we define $E_i := E^s(p_i)$. Since the stable foliation $W_i^s(x) = W_{loc}^s(x) \cap P_i$ of F restricted to $x \in P_i$ is C^1 -close to the foliation of P_i by straight lines with direction E_i when the diameter of the Markov partition is small, we can assume, up to performing a C^1 -perturbation of the unimodal family $f(x, y)$, that the stable foliation of F restricted to P_i is the foliation by straight lines parallel to E_i . Indeed, if we consider the horizontal segments γ_i^u forming the bottom unstable boundary of P_i and we denote by $\{\Pi_i(x, y)\} = ((x, y) + \mathbb{R}E_i) \cap \gamma_i^u$, we can define $g(x, y)$ as the x -coordinate of the point $\{(g(x, y), K_{sgn(x)}(y))\} = (F(\Pi_i(x, y)) + \mathbb{R}E_j) \cap (\mathbb{R} \times \{K_{sgn(x)}(y)\})$ when $F(\Pi_i(x, y)) \in P_j$ and a map $G(x, y) =$

$(g(x, y), K_{sgn(x)}(y))$. Geometrically, G sends the point $\{(x, y)\} = (\Pi_i(x, y) + \mathbb{R}E_i) \cap (\mathbb{R} \times \{y\})$ to $(F(\Pi_i(x, y)) + \mathbb{R}E_j) \cap (\mathbb{R} \times \{K_{sgn(x)}(y)\})$ and F sends $\{(\bar{x}, y)\} = W_i^s(\Pi_i(x, y)) \cap (\mathbb{R} \times \{y\})$ to $W_i^s(F(\Pi_i(x, y))) \cap (\mathbb{R} \times \{K_{sgn(x)}(y)\})$. Since the stable foliation W_i^s is C^1 to the linear foliation $((x, y) + \mathbb{R}E_i) \cap P_i$, the map G is C^1 -close to F and thus $G \in \mathcal{D}^1$ (i.e., $g(x, y)$ is unimodal in the x -variable) as $F \in \mathcal{Z}_\epsilon$. So, from now on, let us suppose that the stable foliation of F restricted to P_i is the foliation by straight lines parallel to E_i .

Recall that the angle between the stable directions E_i and the unstable (horizontal) directions is uniformly bounded away from zero. In particular, we also have a system of coordinates on each P_i (given by the horizontal foliation and the foliation by lines parallel to E_i) where we can write $F|_{P_i}(x, y) = (f_i(x), K_{sgn(x)}(y))$ and $\widetilde{\Lambda}_{\epsilon/2} \cap P_i$ is a product of two dynamically defined Cantor sets, i.e., $\widetilde{\Lambda}_{\epsilon/2} \cap P_i = \mathcal{K}_i^s \cdot (1, 0) + \mathcal{K}_i^u \cdot (\mu_i, 1)$ with $\mathcal{K}_i^s, \mathcal{K}_i^u$ dynamical Cantor sets of the real line and $(\mu_i, 1) \in E_i$.

In this context, the fact that the verticals $\{\pm\epsilon\} \times [0, 1]$ don't intersect $\widetilde{\Lambda}_{\epsilon/2}$ is equivalent to $\pm\epsilon \notin \mathcal{K}_i^s + \mu_i \cdot \mathcal{K}_i^u$ for every $i = 1, \dots, M$. However, this property can be achieved by a C^1 -typical perturbation \widehat{F} of $F \in \mathcal{D}^2$: by Moreira's theorem 4.3, we can choose, for each i , a $(\widehat{\mathcal{K}}_i^s, \widehat{f}_i)$ C^1 -dynamically defined Cantor set C^1 -close to (\mathcal{K}_i^s, f_i) so that $\pm\epsilon \notin \widehat{\mathcal{K}}_i^s + \mu_i \cdot \mathcal{K}_i^u$, and, consequently, $\widehat{F}|_{P_i}(x, y) := (\widehat{f}_i(x), K_{sgn(x)}(y)) \in \mathcal{D}^2$ has the desired property. \square

4.2. (Quasi) Critical points eventually return.

Definition 9. Given $\epsilon > 0$, we call any point $(\pm\epsilon, y)$ with $y \in \mathcal{K}_0$ a ϵ -quasi-critical point (or simply quasi-critical point).

Now we use again C. G. Moreira's fundamental result (Theorem 4.3) to show that, for a C^1 generic $F \in \mathcal{D}^2$, any quasi critical point returns to the "critical region". In other words, roughly speaking, the next result states that we can avoid in the C^1 topology the thickness obstruction (responsible for C^2 Newhouse phenomena).

Lemma 4.3. Let $\epsilon > 0$ be a positive constant. Then, for F in a C^1 -generic (G_δ -dense) subset of \mathcal{D}^1 , there exists $m_0 \in \mathbb{N}$ such that any quasi-critical point $(\pm\epsilon, y) \in \{\pm\epsilon\} \times \mathcal{K}_0$ satisfies

$$F^{m_y}(\pm\epsilon, y) \in R_\epsilon := (-\epsilon, \epsilon) \times [0, 1] \text{ and } F^m(\pm\epsilon, y) \notin (-\epsilon, \epsilon) \times [0, 1], \forall 0 < m < m_y$$

for some positive integer $m_y \leq m_0$ or it is contained in the basin of attraction of some of the (finitely many) attracting periodic points of Theorem 4.1.

Proof. Take $F \in \mathcal{D}^1$ with the properties described during the proof of Corollary 4.1. Since the maximal invariant set Λ_ϵ of U_ϵ is the union of a finite number of periodic sinks and a hyperbolic set $\widehat{\Lambda}_\epsilon$ of saddle type, we see that our task is equivalent to show that

$$\left(\bigcup_{k \geq 0} F^k(\{\pm\epsilon\} \times \mathcal{K}_0) \right) \cap W_{loc}^s(\widehat{\Lambda}_\epsilon) = \emptyset$$

for a C^1 -generic $F \in \mathcal{D}^1$. Keeping this goal in mind, given $N \in \mathbb{N}$, we define

$$\mathcal{G}_N := \left\{ F \in \mathcal{D}^1 : \left(\bigcup_{k=0}^N F^k(\{\pm\epsilon\} \times \mathcal{K}_0) \right) \cap W_{loc}^s(\widehat{\Lambda}_\epsilon) = \emptyset \right\}.$$

It follows that the proof of the lemma is complete once we show that \mathcal{G}_N is C^1 -dense (because it is clearly C^1 -open). Observe that \mathcal{G}_0 is C^1 dense because Λ_ϵ is locally maximal with isolating

neighborhood U_ϵ for a C^1 -typical F (in view of Corollary 4.1). Assuming that \mathcal{G}_{N-1} is C^1 -dense for some $N \geq 1$, we claim that \mathcal{G}_N is also C^1 -dense. In fact, given $F \in \mathcal{G}_{N-1} \cap \mathcal{D}^2$ with the properties appearing in the proof of Corollary 4.1, we can refine the Markov partition $\{P_i\}_{i=1}^M$ so that $F^j(\{\pm\epsilon\} \times \mathcal{K}_0) \cap P_i = \emptyset$ for every $0 \leq j \leq N-1$.

Next, for every $p \in \{\pm\epsilon\} \times [0, 1]$, we denote by $E(p)$ the tangent line of the C^2 curve $F^N(\{\pm\epsilon\} \times [0, 1])$ at the point $F^N(p)$. Note that $E(p)$ is a C^1 function of $p \in \{\pm\epsilon\} \times [0, 1]$. Therefore, since \mathcal{K}_0 is a C^2 dynamical Cantor set of Hausdorff dimension $HD(\mathcal{K}_0) < 1$, we see that, without loss of generality, one can assume that the directions E_i of the stable foliations (by parallel straight lines) of $P_i \cap W_{loc}^s(\hat{\Lambda}_\epsilon)$ don't belong to the set of directions $\{E(p) : p \in \{\pm\epsilon\} \times \mathcal{K}_0\}$. Furthermore, by compactness, we can also fix a Markov partition I_1, \dots, I_k of \mathcal{K}_0 of sufficiently small diameter so that the directions E_i are still transversal to the finite collection of C^2 curves $F^N(\{\pm\epsilon\} \times I_l)$ for every $i = 1, \dots, M$ and $l = 1, \dots, k$. At this stage, we write

$$P_i \cap F^N(\{\pm\epsilon\} \times \mathcal{K}_0) = P_i \cap \bigcup_{b=1}^{a(i)} F^N(\{\pm\epsilon\} \times (\mathcal{K}_0 \cap I_{l(b,i)}))$$

for an adequate choice of indices $l(b, i) \in \{1, \dots, k\}$, and we observe that, by transversality, the projection of each $F^N(\{\pm\epsilon\} \times (I_{l(b,i)} \cap \mathcal{K}_0))$ along the direction E_i gives a C^2 dynamical Cantor set $L_{b,i}$. Moreover, we note that $P_i \cap F^N(\{\pm\epsilon\} \times \mathcal{K}_0) \cap W_{loc}^s(\hat{\Lambda}_\epsilon) \neq \emptyset$ if and only if $\mathcal{K}_i^s \cap (\cup_{b=1}^{a(i)} L_{b,i}) \neq \emptyset$ (where \mathcal{K}_i^s is the stable Cantor set introduced during the proof of Corollary 4.1). Using Moreira's theorem 4.3, we obtain $(\tilde{\mathcal{K}}_i^s, \tilde{f}_i)$ dynamical Cantor sets C^1 -close to (\mathcal{K}_i^s, f_i) such that $\tilde{\mathcal{K}}_i^s \cap (\cup_{b=1}^{a(i)} L_{b,i}) = \emptyset$ for every i . It follows that $\tilde{F}|_{P_i(x, y)} := (\tilde{f}_i(x), K_{sgn(x)}y)$ (in the linearizing coordinates inside each P_i) is C^1 -close to $F \in \mathcal{G}_{N-1}$ and $\tilde{\Lambda}_\epsilon \cap P_i = \tilde{\mathcal{K}}_i^s \times \mathcal{K}_0$ (in the same linearizing coordinates). In particular, by construction, we have $\tilde{F} \in \mathcal{G}_N$. This ends the argument. \square

Remark 4.1. *In the previous statement, we deal with the returns to the critical strip of “quasi-critical” points $(\pm\epsilon, y)$, $y \in \mathcal{K}_0$, instead of critical points c_y^\pm . The technical reason behind this procedure will be clear in the next section (when we perform the “flatness” perturbation to force critical points to fall into the basins of sinks).*

4.3. Creating sinks whose large basins contain all critical points.

Lemma 4.4. *For a C^1 -open and dense subset of $F \in \mathcal{D}^1$, the critical points $c_y^\pm \in \{0^\pm\} \times \mathcal{K}_0$ belong to the union of the basins of a finite number of periodic sinks of F .*

Proof. Fix $F \in \mathcal{D}^1$ be a C^1 -generic map satisfying the properties of Lemma 4.3. Given $\delta > 0$, we will find a C^1 -perturbation of F with size δ whose critical points belong to the basins of finitely many periodic sinks. In this direction, we take $\epsilon > 0$ sufficiently small such that $|\partial_x f(x, y)| < \delta/2$ for every $|x| \leq \epsilon$ and $y \in [0, 1]$. Now, we perturb F to make it “flat” in the critical strip $\overline{R_\epsilon} := [-\epsilon, \epsilon] \times [0, 1]$, i.e., we define

$$g(x, y) = \begin{cases} f(x, y) & \text{if } |x| \geq \epsilon \\ f(\pm\epsilon, y) & \text{if } |x| \leq \epsilon \end{cases}$$

and $G(x, y) := (g(x, y), K_{sgn(x)}(y))$. Observe that, although $G \notin \mathcal{D}^1$ because $g(x, y)$ is not C^1 , G is $\delta/2$ -close to F in the Lipschitz norm and $G = F$ outside the critical strip R_ϵ . In particular, the pieces of orbits of F and G are equal while they stay outside R_ϵ . Hence, since F satisfies Lemma 4.3, we have that G satisfies the same properties, namely, either its quasi-critical points

$\{\pm\epsilon\} \times \mathcal{K}_0$ return to the critical region R_ϵ (after a bounded number of iterates) or they fall into the basins of finitely many periodic sinks (inside Λ_ϵ). We claim that the quasi-critical points returning to R_ϵ belong to the basins of finitely many periodic *sinks* of G . Indeed, by compactness and continuity, we can take a Markov partition I_1, \dots, I_k of K_0 of small diameter and some integers r_1, \dots, r_k so that every quasi-critical point $p \in \{\pm\epsilon\} \times I_l$ return to R_ϵ or fall into the basin of a sink after exactly r_l iterates. Since the pieces of orbits of F and G outside R_ϵ are the same, and the piece of the G -orbit outside R_ϵ of a point $(x, y) \in R_\epsilon$ equals to the piece of F -orbit outside R_ϵ of the point $(\pm\epsilon, y)$, we obtain that G send the boxes $[-\epsilon, \epsilon] \times I_l$ strictly inside another (*a priori* different box) $[-\epsilon, \epsilon] \times I_j$ or inside the basin of a periodic sink after r_l iterates (exactly), so that our claim follows. Finally, we complete the proof by noticing that, although $G \notin \mathcal{D}^1$, one can *slightly* “undo” the “flat” perturbation in order to get a $H \in \mathcal{D}^1$ such that its critical points belong to the basin of finitely many periodic sinks and H is $\delta/2$ -close to G in the Lipschitz norm (and, *a fortiori*, $H \in \mathcal{D}^1$ is δ -close to $F \in \mathcal{D}^1$ in the C^1 -topology). \square

4.4. End of the proof of Theorem B. By combining Corollary 4.1 and Lemma 4.4, we get that the non-wandering set $\Omega(F)$ of a C^1 -typical $F \in \mathcal{D}^2$ can be written as $\Omega(F) = \Lambda_\epsilon \cup \{p_1, \dots, p_k\}$ where p_1, \dots, p_k are periodic sinks of F whose (large) basins contain a ϵ -neighborhood of the critical set, i.e., $\Omega(F)$ is a hyperbolic set.

Thus, the proof of Theorem B will be complete once we can show the following claim: a Kupka-Smale⁶ $F \in \mathcal{D}^1$ such that $\Omega(F)$ is hyperbolic is Axiom A. However, this is a consequence of the following argument of Pujals and Sambarino [PS1, p. 966]: $\Omega(F)$ hyperbolic implies $L(F)$ hyperbolic, so that the results of Newhouse [N4] say that (1) periodic points are dense in $L(F)$ and we can do spectral decomposition of $L(F)$ into finitely many basic sets L_1, \dots, L_k , and (2) we have $\Omega(F) = L(F)$ whenever there is no cycle between the L_i . Hence, we can show that $\Omega(F) = L(F)$ whenever we can verify the no-cycles condition. Since our phase space is two-dimensional, a cycle can only occur among basic sets of saddle-type. However, since F is Kupka-Smale, the intersections of invariant manifolds involved in this cycle are transversal, a contradiction.

REFERENCES

- [ABCD] F. Abdenur, C. Bonatti, S. Crovisier, L. Diaz, Generic diffeomorphisms on compact surfaces, *Fund. Math.*, **187** (2005), 127–159.
- [A] D. Anosov, Extension of zero-dimensional hyperbolic sets to locally maximal ones, *Sb. Math.*, **201** (2010), 935–946.
- [BC] M. Benedicks, L. Carleson, The dynamics of the Hénon map, *Ann. of Math.*, **133** (1991), 73–169.
- [dMvS] W. de Melo, S. van Strien, One-dimensional dynamics, *Springer Verlag*, Berlin, 1993. xiv+605 pp. ISBN: 3-540-56412-8.
- [J] M. Jakobson, Smooth mappings of the circle into itself, *Mat. Sb. (N.S.)*, **85 (127)** (1971), 163–188.
- [M] C. G. Moreira, There are no C^1 -stable intersections of regular Cantor sets, *Acta Math.*, **206** (2011), 311–323.

⁶In principle, we can not directly apply Kupka-Smale theorem in our context because \mathcal{D}^1 is a specific parameter space. However, it is not hard to see that the arguments in the proof of Kupka-Smale theorem are still valid in our case. Indeed, the hyperbolicity of periodic points of a C^1 -generic $F \in \mathcal{D}^1$ was verified in footnote 3, so it remains to check the generic transversality of invariant manifolds of periodic points, but this last task is “automatic” in our setting: we showed that the non-wandering set $\Omega(F)$ of a C^1 -generic $F \in \mathcal{D}^1$ is the union of a hyperbolic set away from the critical line and a finite number of sinks whose basins cover the critical line. In particular, any point q in the intersection of the stable and unstable manifolds of a hyperbolic periodic point p of saddle-type does not approach the critical line and thus the horizontal direction $\mathbb{R} \cdot (1, 0)$ is invariant along the orbit of q . Hence, if the stable and unstable manifolds of p meet non-transversely at q , we would deduce that the stable direction of p is horizontal, a contradiction (as p has saddle-type, so that the horizontal direction is unstable).

- [M1] R. Mañé, Hyperbolicity, sinks and measure in one-dimensional dynamics, *Comm. Math. Phys.*, **100** (1985), 495-524.
- [M2] R. Mañé, Ergodic Theory and Differentiable Dynamics, *Springer Verlag*, Berlin, 1987. xii+317 pp. ISBN: 3-540-15278-4
- [N1] S. Newhouse, Non-density of Axiom A(a) on S^2 , *Proc. A.M.S. Symp. Pure Math.*, **14** (1970), 191-202.
- [N2] S. Newhouse, Diffeomorphism with infinitely many sinks, *Topology*, **13** (1974), 9-18.
- [N3] S. Newhouse, The abundance of wild hyperbolic sets and nonsmooth stable sets for diffeomorphisms, *Publ. Math. Inst. Hautes Études Sci.*, **50** (1979), 101-151.
- [N4] S. Newhouse, Hyperbolic limit sets, *Trans. A.M.S.*, **167** (1972), 125-150.
- [PT] J. Palis, F. Takens, Hyperbolicity and sensitive chaotic dynamics at homoclinic bifurcations, *Cambridge Studies in Adv. Math.*, **35** Cambridge Univ. Press (1993).
- [Pl] V. A. Pliss, On a conjecture due to Smale, *Differencial'nye Uravnenija*, **8** (1972), 268-282.
- [PRH] E. Pujals, F. Rodriguez-Hertz, Critical points for surface diffeomorphisms, *J. Mod. Dyn.*, **4** (2007), 615-648.
- [PS1] E. Pujals, M. Sambarino, Homoclinic tangencies and hyperbolicity for surface diffeomorphisms, *Ann. of Math.*, **151** (2000), no. 3, 961-1023.
- [Sm] S. Smale, Differentiable dynamical systems, *Bull. Amer. Math. Soc.*, **73** (1967), 747-817.
- [U] R. Ures, Abundance of hyperbolicity in the C^1 topology, *Ann. Sci. Éc. Norm. Sup. (4)*, **28** (1995), no. 6, 747-760.
- [U2] R. Ures, On the approximation of Hénon-like attractors by homoclinic tangencies, *Ergodic Theory Dynam. Systems*, **15** (1995), 1223-1229.

CARLOS MATHEUS: UNIVERSITÉ PARIS 13, SORBONNE PARIS CITÉ, LAGA, CNRS (UMR 7539), F-93439, VILLETANEUSE, FRANCE.

E-mail address: matheus@impa.br.

URL: <http://www.impa.br/~cmateus>

CARLOS G. MOREIRA: INSTITUTO DE MATEMÁTICA PURA E APLICADA (IMPA), ESTRADA D. CASTORINA, 110, J. BOTÂNICO, 22460-320, RIO DE JANEIRO, BRAZIL.

E-mail address: gugu@impa.br.

URL: <http://www.impa.br/~gugu>

ENRIQUE PUJALS: INSTITUTO DE MATEMÁTICA PURA E APLICADA (IMPA), ESTRADA D. CASTORINA, 110, J. BOTÂNICO, 22460-320, RIO DE JANEIRO, BRAZIL.

E-mail address: enrique@impa.br.

URL: <http://www.impa.br/~enrique>