

# Dynamics: a long century

Marcelo Viana

IMPA - Rio de Janeiro

## The $n$ -body problem

$$\frac{d^2 \vec{r}_i}{dt^2} = \sum_{i \neq j} G m_i m_j \frac{\vec{r}_j - \vec{r}_i}{\|\vec{r}_j - \vec{r}_i\|^3} \quad i = 1, 2, \dots, n$$

can not solved analytically, in general, when  $n > 2$ .

## The $n$ -body problem

$$\frac{d^2 \vec{r}_i}{dt^2} = \sum_{i \neq j} G m_i m_j \frac{\vec{r}_j - \vec{r}_i}{\|\vec{r}_j - \vec{r}_i\|^3} \quad i = 1, 2, \dots, n$$

can not solved analytically, in general, when  $n > 2$ .


Poincaré proposes to use tools from geometry, probability, analysis, algebra, to describe the qualitative behavior of solutions without actually computing them.

## Continuous time and discrete time

Flow (associated to a vector field  $F$ )

$\{f^t : M \rightarrow M : t \in \mathbb{R}\}$  defined by  $f^t(a) =$  value at time  $t$  of the solution to

$$\frac{dx}{dt} = F(x) \text{ and } x(0) = a.$$




## Continuous time and discrete time

Flow (associated to a vector field  $F$ )

$\{f^t : M \rightarrow M : t \in \mathbb{R}\}$  defined by  $f^t(a) =$  value at time  $t$  of the solution to

$$\frac{dx}{dt} = F(x) \text{ and } x(0) = a.$$



Iteration of an (invertible) transformation

$f : M \rightarrow M$  invertible map; one denotes by  $f^n$  the  $n$ -fold iterate.


## Return maps and suspensions

One can relate continuous time systems and discrete time systems:

## Return maps and suspensions

One can relate continuous time systems and discrete time systems:

continuous  $\rightarrow$  discrete




$f : \Sigma' \rightarrow \Sigma$  is the **first return map** of the flow to the cross-section.

## Return maps and suspensions

One can relate continuous time systems and discrete time systems:

continuous  $\rightarrow$  discrete



$f : \Sigma' \rightarrow \Sigma$  is the **first return map** of the flow to the cross-section.

continuous  $\leftarrow$  discrete

Every invertible transformation  $f$  can be realized as a first return map of some flow, that we call **suspension flow** of  $f$ .



## Poincaré recurrence


### Theorem (Poincaré)

*If  $f : M \rightarrow M$  preserves a probability  $\mu$  then, given any measurable set  $B \subset M$ , the orbit of  $\mu$ -almost any  $x \in B$  returns to  $B$ .*

# Poincaré recurrence

## Theorem (Poincaré)

*If  $f : M \rightarrow M$  preserves a probability  $\mu$  then, given any measurable set  $B \subset M$ , the orbit of  $\mu$ -almost any  $x \in B$  returns to  $B$ .*




Let  $E = \{\text{points whose orbits never return to } B\}$ . Then  $\mu(E) = 0$ :

$$\sum_n \mu(E) = \sum_n \mu(f^n(E)) = \mu\left(\bigcup_n f^n(E)\right) \leq \mu(M) = 1.$$

# Twist maps

## Theorem (Poincaré-Birkhoff)


*Assume  $f : A \rightarrow A$  preserves the boundary and the area measure and twists the radii. Then there is  $p \in A$  such that  $f(p) = p$ .*



## Transverse homoclinic intersections

### Theorem (Birkhoff)

*Any transverse homoclinic point  $q$  associated to a saddle point  $p$  is accumulated by periodic points,  $f^{k_i}(p_i) = p_i$ , with periods  $k_i \rightarrow \infty$ .*



# The ergodic hypothesis

**Boltzmann:** In the long run, a system of molecules will assume all conceivable micro-states that are compatible with the conservation of energy.

# The ergodic hypothesis

**Boltzmann:** In the long run, a system of molecules will assume all conceivable micro-states that are compatible with the conservation of energy.

Reformulation: Over long periods of time, the time spent by the system in some region of the phase space of microstates with the same energy is proportional to the volume of that region.

In other words, all micro-states with the same energy should be equally probable, in the long run. Is this so ?

# The ergodic theorem

## Theorem (von Neumann, Birkhoff)

Assume  $f : M \rightarrow M$  preserves a probability  $\mu$ . Then, for any integrable function  $\psi : M \rightarrow \mathbb{R}$ , the time-average

$$\tilde{\psi}(x) = \lim_{n \rightarrow \infty} \frac{1}{n} (\psi(f(x)) + \cdots + \psi(f^n(x)))$$

exists for  $\mu$ -almost every point  $x$ .

# The ergodic theorem

## Theorem (von Neumann, Birkhoff)

Assume  $f : M \rightarrow M$  preserves a probability  $\mu$ . Then, for any integrable function  $\psi : M \rightarrow \mathbb{R}$ , the time-average

$$\tilde{\psi}(x) = \lim_{n \rightarrow \infty} \frac{1}{n} (\psi(f(x)) + \cdots + \psi(f^n(x)))$$

exists for  $\mu$ -almost every point  $x$ .

If one takes  $\psi =$  characteristic function of some set  $B \subset M$ , then  $\tilde{\psi}(x)$  is the time the orbit of  $x$  spends in the set  $B$ .



# The ergodic theorem

## Theorem (von Neumann, Birkhoff)

Assume  $f : M \rightarrow M$  preserves a probability  $\mu$ . Then, for any integrable function  $\psi : M \rightarrow \mathbb{R}$ , the time-average


$$\tilde{\psi}(x) = \lim_{n \rightarrow \infty} \frac{1}{n} (\psi(f(x)) + \cdots + \psi(f^n(x)))$$

exists for  $\mu$ -almost every point  $x$ .

If one takes  $\psi =$  characteristic function of some set  $B \subset M$ , then  $\tilde{\psi}(x)$  is the time the orbit of  $x$  spends in the set  $B$ .

The system is **ergodic** if the time-averages are constant  $\mu$ -almost everywhere.

## Geodesic flows on surfaces



### Theorem (Hedlund, Hopf)

*The geodesic flow on a surface with negative curvature is ergodic, relative to the Liouville measure.*

# KAM theory

## Theorem (Kolmogorov, Arnold, Moser)


*If  $f : A \rightarrow A$  is a  $C^\infty$  area preserving twist map, there exists a subset of  $A$  with positive area consisting of closed curves that are fixed by  $f$ . In particular,  $f$  is **not** ergodic.*

This is a manifestation of a very general phenomenon that applies, in particular, to volume preserving (symplectic) transformations and flows in any dimension.

# The horseshoe

## Theorem (Smale)


*If  $f$  has some transverse homoclinic point then some iterate  $f^N$  exhibits a horseshoe.*



# The horseshoe

## Theorem (Smale)

*If  $f$  has some transverse homoclinic point then some iterate  $f^N$  exhibits a horseshoe.*



So,  $f$  has periodic points with arbitrarily large periods (Birkhoff). This explains how an infinite number of periodic points can coexist in a robust fashion.

# Hyperbolicity

The **limit set** of  $f : M \rightarrow M$  is the closure  $L(f)$  of the set of all accumulation points of all orbits (forward and backward).

# Hyperbolicity

The **limit set** of  $f : M \rightarrow M$  is the closure  $L(f)$  of the set of all accumulation points of all orbits (forward and backward).

The transformation is **hyperbolic** if the tangent space admits a splitting

$$T_x M = E_x^s \oplus E_x^u$$

at every point  $x \in L(f)$ , invariant under the derivative and such that  $Df$  contracts  $E^s$  and  $Df^{-1}$  contracts  $E^u$ , at uniform rates.

# Hyperbolicity

The **limit set** of  $f : M \rightarrow M$  is the closure  $L(f)$  of the set of all accumulation points of all orbits (forward and backward).

The transformation is **hyperbolic** if the tangent space admits a splitting

$$T_x M = E_x^s \oplus E_x^u$$

at every point  $x \in L(f)$ , invariant under the derivative and such that  $Df$  contracts  $E^s$  and  $Df^{-1}$  contracts  $E^u$ , at uniform rates.

When  $L(f) = M$ , we say  $f$  is **globally hyperbolic** or **Anosov**.



# Geodesic flows on manifolds

## Theorem (Anosov)

- 1 *The geodesic flow on any manifold with negative sectional curvature is globally hyperbolic (and preserves the Liouville volume measure).*
- 2 *Every globally hyperbolic system that preserves the volume measure is ergodic.*

## Dynamical decomposition

### Theorem (Smale, Newhouse)

*If  $f$  is hyperbolic then  $L(f) = \Lambda_1 \cup \dots \cup \Lambda_N$  where the  $\Lambda_i$  are invariant, disjoint, and indecomposable (contain dense orbits). The forward/backward accumulation set of the orbit of every  $x \in M$  is contained in some  $\Lambda_i$ .*

## Dynamical decomposition

### Theorem (Smale, Newhouse)

*If  $f$  is hyperbolic then  $L(f) = \Lambda_1 \cup \dots \cup \Lambda_N$  where the  $\Lambda_i$  are invariant, disjoint, and indecomposable (contain dense orbits). The forward/backward accumulation set of the orbit of every  $x \in M$  is contained in some  $\Lambda_i$ .*

$\Lambda_i$  is an **attractor** if the **basin of attraction**

$$B(\Lambda_i) = \{x \in M \text{ whose forward orbit accumulates in } \Lambda_i\}$$

has positive volume (then it is a neighborhood of  $\Lambda_i$ ).

The basins of the attractors of a hyperbolic transformation cover a full volume set.

## Physical measures

### Theorem (Sinai, Ruelle, Bowen)

*Given any hyperbolic attractor  $\Lambda$ , there exists a probability  $\mu_{SRB}$  supported on  $\Lambda$  such that every time-average*

$$\lim_n (\psi(f(x)) + \cdots + \psi(f^n(x))) = \int \psi d\mu_{SRB}$$

*at volume-almost every point  $x$  in the basin of attraction of  $\Lambda$ .*

## Beyond hyperbolicity ?



However, many important dynamical systems are **not** hyperbolic.

To what extent can we develop a similarly rich theory for very general dynamical systems?

*Dynamics beyond uniform hyperbolicity*, Bonatti, Díaz, Viana,  
Springer Verlag.


# Homoclinic tangencies

Only two mechanisms are known that yield robustly non-hyperbolic behavior:



# Heterodimensional cycles

Only two mechanisms are known that yield robustly non-hyperbolic behavior:



## Strong density conjecture


### Conjecture (Palis)

Every dynamical system can be approximated by one which either is hyperbolic, has a homoclinic tangency, or a heterodimensional cycle.



# Hénon strange attractor

The **Hénon map** is defined by  $f_{a,b}(x, y) = (1 - ax^2 + by, x)$  where  $a$  and  $b$  are parameters. Hénon observed that it seems to have a “strange” attractor:



## Abundance of strange attractors

### Theorem (Benedicks, Carleson, Young, Viana)

*For a positive measure set of parameters  $a$  and  $b$ ,*

**BC** *the Hénon map has a strange (non-hyperbolic) attractor  $\Lambda$*

**BY** *the attractor  $\Lambda$  supports some physical measure  $\mu_{phys}$*

**BV**  *$\tilde{\psi}(x) = \int \psi d\mu_{phys}$  for volume-almost all  $x \in B(\Lambda)$  and all  $\psi$*

**BV** *the dynamics in the basin is stable under random noise.*

## Abundance of strange attractors

### Theorem (Benedicks, Carleson, Young, Viana)

*For a positive measure set of parameters  $a$  and  $b$ ,*

**BC** *the Hénon map has a strange (non-hyperbolic) attractor  $\Lambda$*

**BY** *the attractor  $\Lambda$  supports some physical measure  $\mu_{phys}$*

**BV**  *$\tilde{\psi}(x) = \int \psi d\mu_{phys}$  for volume-almost all  $x \in B(\Lambda)$  and all  $\psi$*

**BV** *the dynamics in the basin is stable under random noise.*

### Theorem (Mora, Viana)


*Hénon-like strange attractors occur, in a persistent way, whenever a dissipative homoclinic tangency is unfolded.*

There is considerable recent progress by Wang, Young.

## Lorenz strange attractors

The **Lorenz flow** is defined by the system of differential equations

$$\begin{aligned}\dot{x} &= -\sigma x + \sigma y & \sigma &= 10 \\ \dot{y} &= \rho x - y - xz & \rho &= 28 \\ \dot{z} &= xy - \beta z & \beta &= 8/3\end{aligned}$$



## Robust strange attractors

### Theorem (Morales, Pujals, Pacifico)

*Every robust attractor  $\Lambda$  of a flow in 3 dimensions is either hyperbolic or Lorenz-like (singular-hyperbolic): the latter occurs precisely if the attractor contains some equilibrium point.*

**Robust** means that there exists a neighborhood  $U$  of  $\Lambda$  such that the set of points whose orbits never leave  $U$

- is indecomposable (dense orbits), for any perturbation of the original flow
- and coincides with  $\Lambda$  for the original flow.

## Robust strange attractors

### Theorem (Morales, Pujals, Pacifico)

*Every robust attractor  $\Lambda$  of a flow in 3 dimensions is either hyperbolic or Lorenz-like (singular-hyperbolic): the latter occurs precisely if the attractor contains some equilibrium point.*

**Robust** means that there exists a neighborhood  $U$  of  $\Lambda$  such that the set of points whose orbits never leave  $U$

- is indecomposable (dense orbits), for any perturbation of the original flow
- and coincides with  $\Lambda$  for the original flow.

### Theorem (Tucker)

*The Lorenz original equations do exhibit a strange attractor.*

## Partial hyperbolicity

A transformation is **partially hyperbolic** if the tangent space admits an invariant splitting

$$T_x = E_x^s \oplus E_x^c \oplus E_x^u$$

at every point, such that  $Df | E^s$  is a contraction,  $Df | E^u$  is an expansion, and  $Df | E^c$  is “in between” the two, with uniform rates.

## Partial hyperbolicity

A transformation is **partially hyperbolic** if the tangent space admits an invariant splitting

$$T_x = E_x^s \oplus E_x^c \oplus E_x^u$$

at every point, such that  $Df|E^s$  is a contraction,  $Df|E^u$  is an expansion, and  $Df|E^c$  is “in between” the two, with uniform rates.

This notion turns out to be crucial for understanding

- **robust indecomposability**: Bonatti, Díaz, Pujals, Ures, Viana, Arbieto, Matheus, Horita, Tahzibi
- **stable ergodicity**: Grayson, Pugh, Shub, Wilkinson, Burns, Dolgopyat, Nitika, Torok, Xia, Bonatti, Matheus, Viana, Rodriguez-Herz (F & J), Ures



## Two conjectures

### Finiteness of attractors (Palis)

Every dynamical system is approximated by one having only finitely many attractors, and these attractors support physical measures.

## Two conjectures

### Finiteness of attractors (Palis)

Every dynamical system is approximated by one having only finitely many attractors, and these attractors support physical measures.

### Hyperbolicity and physical measures (Viana)

If there is an invariant splitting  $T_x M = E_x^1 \oplus E_x^2$  at volume-almost every point, such that  $Df^n | E^1$  is eventually contracting, and  $Df^n | E^2$  is eventually expanding, then the system admits some physical measure.